



Supporting Next Century's Science

Biochemical Reagents

6th
Edition

Contents

Nucleic Acid Isolation/ Electrophoresis

Total RNA Isolation Reagent: Sepasol-RNA I Super G	4
Tail Lysis Buffer	6
Nuclease Decontamination: RNase Quiet	7
Coprecipitant of DNA/RNA: Gene-Packman Coprecipitant	8
Agarose for Nucleic Acid Electrophoresis	9
Bullet PAGE Plus Precast Gel	11
DNA Ladder Markers	12
Dual Green Nucleic Acid Stain.....	13
Ethidium Bromide Solution (0.44 mg/ml)	14
Phenol solution for DNA/RNA extraction	15

Cell Culture

Cell Culture Medium	16
Custom Cell Culture Media	18
Reduced Serum DMEM/Ham's F-12 Medium	18
AscleStem series	20
AscleStem [®] hPSC Maintenance Supplement.....	21
AscleStem [®] Dissociation Solution for hPSCs	22
AscleStem [®] Hepatocyte Differentiation & Maintenance Basal Medium	23
AscleStem [®] Neuronal Medium & Supplement(50x)	24
AscleStem [®] 0.1%-Gelatin Solution	26
0.01%-Polyethyleneimine Coating Solution.....	27
Mitomycin C Solution (1 mg/ml) for preparation of feeder cells	27
Cell Dissociation Reagent: Accutase [™]	28
EZSPHERE [®] for 3D cell culture with microwells.....	29
Serum-free Cell Freezing Media: Cell Reservoir One	30
ES/iPS cell Freezing Media: Cell Reservoir One, Vitrify	31
Balanced Salines	32
Sterile water	32
Media additives	33
Reagents for cell dissociation	36
Antibiotics	36
Cell Count Reagent SF, based on WST-8	37
MTT Cell Count Kit, based on reduction of MTT.....	38
LDH Cytotoxicity Assay Kit	38
Trypan Blue Solution.....	39
Reagents for apoptosis research.....	39
Reagents for alkaline phosphatase staining	39
Medium for Bacteria, Plusgrow II.....	40
IPTG and X-gal Solutions	41
Gelling Agent for Plant Study: Gellan Gum.....	42

Cell Extraction / Protein Assay

Zymolyase [™] (from <i>Arthrobacter Luteus</i>)	43
Cell Lysis Solution: Cell Lysis Buffer	44
Protease Inhibitor Cocktail	46
Phosphatase Inhibitor Cocktail	47
Determination of Protein Concentration: Protein Assay.....	48
Protein Assay CBB Clean Up Kit	49
Reductant adaptable reagent for protein assay BCA Kit	50

Contents

Protein Purification

COSMOGEL [®] Ig-Accept Protein G	51
COSMOGEL [®] GST-Accept	52
High Performance Magnetic Nanoparticles: FG beads [®]	53
TurboTEV Protease & Turbo3C Protease.....	54

Protein Electrophoresis


Bullet PAGE Plus Precast Gel	55
Extra PAGE One Precast Gel	59
Electrophoresis Tank for Precast Gel	61
WIDE RANGE Gel Preparation Buffer (4x) for PAGE.....	61
Stacking Gel Buffer Solution (4x) with Blue Color	62
Rapid Running Buffer Solution	63
Polyacrylamide Gel Casting Reagents	64
Running Buffers	64
Sample Buffer Solution for SDS-PAGE (6x)	65
Molecular Weight Markers	66
PAGE Clean Up Kit.....	68
CBB Stain Solution	69
Silver Staining Kit.....	71

Western Blotting

Bullet Semi-dry Transfer One	73
Blocking One and Blocking One P.....	75
Bullet Blocking One for Western Blotting.....	76
Bullet ImmunoReaction Buffer	79
Signal Enhancer HIKARI for Western blotting and ELISA	80
HRP-conjugated secondary antibody	81
Chemiluminescent Western Blotting Substrates.....	82
Colorimetric Western Blotting Substrates	84
WB Stripping Solution.....	86
Epitope Tag Antibody	87

Immunohistochemistry

4% - Paraformaldehyde Phosphate Buffer Solution	89
HistoVT One (10x, pH 7.0)	90
Blocking One Histo	91
Signal Enhancer HIKARI for Immunostain.....	92
Anti-GFP (Rat IgG2a), Monoclonal (GF090R)	93
Streptavidin Biotin Complex Peroxidase Kit	93
High Sensitivity Peroxidase DAB Stain.....	94
Mounting Medium for Fluorescent Staining	96



Total RNA Isolation Reagent: Sepasol[®]-RNA I Super G

- » Easy to identify interphase
- » Less than 1hr for RNA isolation
- » Enables to extract DNA and proteins along with RNA from a single sample



Phenol Phase Color

Sepasol-RNA I Super G


Aqueous phase

Phenol phase



Easy to identify interphase

Procedure





For details, refer to the instruction manual.

Performance evaluation

Total RNA was extracted from HL-60 cells at 8×10^6 cells, 1×10^6 cells, and 0.125×10^6 cells using this product. Additionally, for the 0.125×10^6 cells of HL-60 cells, samples were prepared with Gene-Packman Coprecipitant (#12680-30) as well as general alcohol precipitation. The extracted total RNA was treated with DNase according to the instruction manual attached and proceed to RT-qPCR. Please note that throughout the entire process, all cell number samples were treated under identical conditions, including the volume of the dissolution solution.

[Result (RT-qPCR)]

Using this product, it was possible to extract mRNA in a cell number-dependent manner. Moreover, although RNA yield decreased and greater variability was observed (yellow lines) when the cell number was low, such as 0.125×10^6 cells, Gene-Packman co-precipitant (#12680-30) increased the yield and reduced variability (orange lines).



Sample	Ct value
8×10^6 cells	13.55 - 13.8
1×10^6 cells	16.12 - 16.35
0.125×10^6 cells	23.04 - 27.87
0.125×10^6 cells (co-precipitation)	19.17 - 19.59
Non-Template Control	34.99

*Threshold: 4.47

(Conditions)

Primer:

[Gene target] β -Actin

[Forward primer] 5'-AAGAGAGGCATCCTCACCT-3'

[Reverse primer] 5'-TACATGGCTGGGGTGTGAA-3'

[PCR product] 218 bp

PCR enzyme (reverse transcriptase enzyme):

One Step TB Green® PrimeScript RT-PCR Kit II

(Perfect Real Time) (Takara Bio #RR086A)

(Thermal Cycler Dice, TB Green and PrimeScript are registered by Takara Bio.)

PCR condition:

1. Hold (reverse transcription) 42°C (5 mins.) → 95°C (10 sec.), 1 cycle

2. 2 Step PCR 95°C (5 sec.) → 60°C (30 sec.), 40 cycles

3. Dissociation 95°C (15 sec.) → 60°C (30 sec.) → 95°C (5 sec.), 1 cycle

Real-time PCR equipment

Thermal Cycler Dice® Real Time System III

Ordering Information

Product Name	Storage	Product No.	PKG Size
Sepasol®-RNA I Super G (for biological samples [cell, tissue, etc.])	R	09379-26	10 ml
		09379-84	100 ml
		09379-97	200 ml
		09379-55	500 ml
Sepasol®-RNA II Super (for liquid samples [blood, etc.])	R	30487-46	100 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Tail Lysis Buffer


Tail Lysis Buffer is ready-to-use solution that enables simple genotyping procedure.

- » Ready-to-use solution
- » DNase, RNase free



Application: Genotyping of mouse tail

Procedure



Sry gene (132 bp), which only exists in male mouse Y-chromosomes was increased in male derived DNA, but was not increased in female derived DNA. The result shows that PCR operates correctly.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Tail Lysis Buffer	RT	06169-95	500 ml

Related product

Product Name	Storage	Product No.	PKG Size
Proteinase K from Tritirachium album	R	29442-14	100 mg
		29442-85	500 mg
Proteinase K(Recombinant) Solution	R	15679-06	2 ml
		15679-64	10 ml


[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Nuclease Decontamination: RNase Quiet

- » **Free from contamination** : removes nucleases and nucleic acids
- » **Easy to wipe** : no detergent contained
- » **Easy to use** : spray type
- » **Non-carcinogenic** : no DEPC contained



Application 1: Decontamination of cover glass




Condition


1. Apply 100 μ l of RNase A solution (1 mg/ml) to cover glasses and dry them.
2. Spray with DEPC-treated water or RNase Quiet and wait for 1 minute. Wipe thoroughly with a clean paper towel and then rinse with RNase-free sterile water.
3. Apply 50 μ l of RNA solution (40 μ g/ml) on the cover glasses and incubate them at 37°C for 30 minutes.
4. Apply 1 μ l ethidium bromide solution (20 μ g/ml) to the cover glasses.
5. Observe with UV detector.

Application 2: RNase and DNase decontamination of 1.5 ml micro-tubes

【RNase decontamination test】



【DNase decontamination test】



Condition

- ① Add the following solutions and dry them up.
 - RNase decontamination test
 - (+): Add 10 μ l of RNase A solution (10 mg/ml) to a 1.5 ml micro-tube.
 - (-): Add 10 μ l of only the solution used to dissolve RNase A to a 1.5 ml micro-tube as a control.
 - DNase decontamination test
 - (+): Add 10 μ l of DNase I (27.3 U/10 μ l) to a 1.5 ml micro-tube, and dry it up.
 - (-): Add 10 μ l of only the solution used to dissolve DNase I to a 1.5 ml micro-tube as a control.
- ② Add 1 ml of DEPC-treated water or RNase Quiet and wait for 1 minute.
- ③ Remove the solution from the tubes and rinse those with 1 ml of DEPC-treated water.
- ④ Remove the DEPC-treated water from the tubes and add 25 μ l of RNA solution (40 μ g/ml) for RNase decontamination test; 25 μ l of DNA solution (40 μ g/ml) and 1 μ l of 50 mmol/L magnesium chloride for DNase decontamination test, and incubate them at 37°C for 30 minutes.
- ⑤ Reacted samples are electrophoresed on agarose gel and detected using EtBr.

Ordering Information

Product Name	Storage	Product No.	PKG Size
RNase Quiet (with spray nozzle)	RT	09147-14	475 ml
RNase Quiet Refill	RT	09477-94	475 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Coprecipitant of DNA/RNA: Gene-Packman Coprecipitant


- » Almost complete nucleic acid recovery
- » No requirement of low-temperature incubation
- » Endotoxin, DNase and RNase tested
- » High nucleic acid pellet visibility

No Effect of Gene-Packman Coprecipitant to PCR

In order to figure out that Gene-Packman Coprecipitant does not effect PCR performance, Gene-Packman Coprecipitant was consciously added to PCR reaction solutions which amplify 1,000 bp DNA fragments. This data shows that Gene-Packman does not affect PCR performance.

Amount of Gene-Packman Coprecipitant to PCR reactin solution:

Lane 1: 0 μ l, Lane 2: 0.2 μ l, Lane 3: 0.5 μ l, Lane 4: 1.0 μ l, Lane 5: 3.0 μ l and Lane 6: 5.0 μ l.



PCR condition

Reaction Volume : 25 μ l reaction

Tempate DNA : 15 ng

Denaturation : 94°C for 0.5 min.

Annealing : 55°C for 0.5 min.

Extension : 72°C for 1 min.

Cycle : 25 times

The agarose gel image above shows no interference with Gene-Packman Coprecipitant to PCR performance.

(Blend Taq®, KOD Dash® and KOD FX® are registered by Toyobo. TaKaRa Ex Taq® are registered by Takara Bio.)

Ordering Information

Product Name	Storage	Product No.	PKG Size
Gene-Packman Coprecipitant	R	12680-30	1 KIT

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Agarose for Nucleic Acid Electrophoresis


● Fine-powdered Agarose

- » **High solubility** : Smaller average particle size for easy dissolution
- » **Simple** : Easy-to-weigh
- » **High sharpness** : Sharp bands observable



Solubility Comparison (Particle size and solubility)

Solubility tests have been done using microwave oven (400W).
Photo image: microscope (x25)



Agarose:	Conventional	Fine-powdered	A Company	B Company
Solubility (speed):	Fast	Very fast	Very fast	Normal
Time:	2 min 50 sec	1 min 53 sec	1 min 55 sec	3 min 10 sec

Specification

Type	: $\geq 1\text{ kbp}$
Sulfate (%)	: ≤ 0.2
Gel Strength	: $\geq 2,500\text{ g/cm}^2$ (at 1.5%)
Gel Point (°C)	: 36 ± 1.5
Electroendosmosis (-mr)	: 0.09-0.13



Ordering Information

Product Name	Storage	Product No.	PKG Size
Agarose for $\geq 1\text{ kbp}$ fragment (Fine Powder)	RT	02468-24	10 g
		02468-66	100 g
		02468-95	500 g

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

● Other Agaroses

Agarose Gel Selection Guide



Ordering Information

Product Name		Storage	Product No.	PKG Size
Agarose-LM Plaque for \geq 1kbp fragment				
Type	: Low Melting Agarose \geq 1kbp fragment	RT	01650-02	25 g
Sulfate (%)	: \leq 0.5		01650-86	100 g
Gel Strength	: \geq 250 g/cm ² (at 1.5%)			
Gel Point (°C)	: \leq 30			
Melting Temp. (°C)	: \leq 65.5			
Electroendosmosis (-mr)	: \leq 0.12			
DNase, RNase tested				
Agarose-LM Sieve for \leq 1kbp fragment				
Type	: Low Melting Agarose \leq 1kbp fragment	RT	01651-92	25 g
Sulfate (%)	: \leq 0.5		01651-76	100 g
Gel Strength	: \geq 1,000 g/cm ² (at 4%)			
Gel Point (°C)	: \leq 35			
Melting Temp. (°C)	: \leq 65			
Electroendosmosis (-mr)	: \leq 0.12			
DNase, RNase tested				
Agarose-RE for \geq 1kbp fragment, for Restriction and Ligation				
Type	: \geq 1kbp	RT	01149-92	25 g
Sulfate (%)	: \leq 0.2		01149-76	100 g
Gel Strength	: \geq 2,500 g/cm ² (at 1.5%)		01149-05	500 g
Gel Point (°C)	: 36 \pm 1.5			
Electroendosmosis (-mr)	: 0.09 - 0.13			
Tested for Silver staining, Dnase, RNase and Enzyme reaction				
Agarose for \geq 1kbp fragment (Fine Powder) refer to pxx				
Type	: \geq 1kbp	RT	02468-24	10 g
Sulfate (%)	: \leq 0.5		02468-66	100 g
Gel Strength	: \geq 2,500 g/cm ² (at 1.5%)		02468-95	500 g
Gel Point (°C)	: 36 \pm 1.5			
Electroendosmosis (-mr)	: 0.09 - 0.13			
Agarose for \geq 1kbp fragment				
Type	: \geq 1kbp	RT	01163-92	25 g
Sulfate (%)	: \leq 0.2		01163-76	100 g
Gel Strength	: \geq 2,500 g/cm ² (at 1.5%)		01163-05	500 g
Gel Point (°C)	: 36 \pm 1.5			
Electroendosmosis (-mr)	: 0.09 - 0.13			
Agarose for 150-1,500bp fragment				
Type	: 150-1,500bp	RT	01153-22	25 g
Sulfate (%)	: \leq 0.1		01153-64	100 g
Gel Strength	: \geq 2,000 g/cm ² (at 1.5%)			
Gel Point (°C)	: \leq 36.5			
Electroendosmosis (-mr)	: \leq 0.12			
DNase, RNase tested				
Agarose for 50-800bp fragment				
Type	: 50-800bp	RT	01147-12	25 g
Sulfate (%)	: \leq 0.1		01147-96	100 g
Gel Strength	: \geq 750 g/cm ² (at 1.5%)			
Gel Point (°C)	: 30			
Electroendosmosis (-mr)	: \leq 0.12			
DNase, RNase tested				
Agarose-LE, Classic type				
Type	: LE, Classic	RT	01157-82	25 g
Sulfate (%)	: \leq 0.2		01157-66	100 g
Gel Strength	: \geq 2,500 g/cm ² (at 1.5%)		01157-95	500 g
Gel Point (°C)	: 36 \pm 1.5			
Electroendosmosis (-mr)	: 0.09 - 0.13			
Agarose-ME, Classic type				
Type	: ME, Classic	RT	01158-72	25 g
Sulfate (%)	: \leq 0.25		01158-56	100 g
Gel Strength	: \geq 2,000 g/cm ² (at 1.5%)		01158-85	500 g
Gel Point (°C)	: 36 \pm 1.5			
Electroendosmosis (-mr)	: 0.16 - 0.19			
Agarose-LM (melting temperature \leq 65°C)				
Type	: Low Melting Agarose	RT	01161-12	25 g
Sulfate (%)	: \leq 0.2		01161-54	100 g
Gel Strength	: \geq 550 g/cm ² (at 1.5%)			
Gel Point (°C)	: 26 \pm 2			
Melting Temp. (°C)	: \leq 65			
Electroendosmosis (-mr)	: \leq 0.12			

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Bullet PAGE Plus Precast Gel

Bullet PAGE Plus is a precast gel for high-speed electrophoresis.

For protein electrophoresis, refer to p54.

» **Only 12 minutes electrophoresis time for nucleic acid with 400 V**

» **Surfactant-free, so can be used for nucleic acid analysis.**

*Please use 25 mM Tris and 192 mM glycine buffer for nucleic acid electrophoresis.

Gel		Gradient gel				
Gel concentration		5-10 %	5-15 %	5-20 %	7.5-15 %	10-20 %
Product number	13 wells	21789-34	21791-84	21793-64	21795-44	21797-24
	17 wells	21790-94	21792-74	21794-54	21796-34	21798-14
Images of nucleic acids (PAGE)		(bp)	(bp)	(bp)	(bp)	(bp)
		1,500→	1,500→	1,500→	1,500→	1,500→
		500→	500→	500→	500→	500→
		100→	100→	100→	100→	100→
		20→	20→	20→	20→	20→

Gel		Homogeneous gel			
Gel concentration		7.5%	10%	12.5%	15%
Product number	13 wells	21799-04	21801-44	21807-84	21853-74
	17 wells	21800-54	21806-94	21811-14	21854-64
Images of nucleic acids (PAGE)		(bp)	(bp)	(bp)	(bp)
		1,500→	1,500→	1,500→	1,500→
		500→	500→	500→	500→
		100→	100→	100→	100→
		20→	20→	20→	20→

Ordering Information

Product Name	Storage	Product No.	PKG Size
Bullet PAGE Plus Precast Gel, 5-10%, 13wells	R	21789-34	10 Sheets
Bullet PAGE Plus Precast Gel, 5-10%, 17wells	R	21790-94	10 Sheets
Bullet PAGE Plus Precast Gel, 5-15%, 13wells	R	21791-84	10 Sheets
Bullet PAGE Plus Precast Gel, 5-15%, 17wells	R	21792-74	10 Sheets
Bullet PAGE Plus Precast Gel, 5-20%, 13wells	R	21793-64	10 Sheets
Bullet PAGE Plus Precast Gel, 5-20%, 17wells	R	21794-54	10 Sheets
Bullet PAGE Plus Precast Gel, 7.5-15%, 13wells	R	21795-44	10 Sheets
Bullet PAGE Plus Precast Gel, 7.5-15%, 17wells	R	21796-34	10 Sheets
Bullet PAGE Plus Precast Gel, 10-20%, 13wells	R	21797-24	10 Sheets
Bullet PAGE Plus Precast Gel, 10-20%, 17wells	R	21798-14	10 Sheets
Bullet PAGE Plus Precast Gel, 7.5%, 13wells	R	21799-04	10 Sheets
Bullet PAGE Plus Precast Gel, 7.5%, 17wells	R	21800-54	10 Sheets
Bullet PAGE Plus Precast Gel, 10%, 13wells	R	21801-44	10 Sheets
Bullet PAGE Plus Precast Gel, 10%, 17wells	R	21806-94	10 Sheets
Bullet PAGE Plus Precast Gel, 12.5%, 13wells	R	21807-84	10 Sheets
Bullet PAGE Plus Precast Gel, 12.5%, 17wells	R	21811-14	10 Sheets
Bullet PAGE Plus Precast Gel, 15%, 13wells	R	21853-74	10 Sheets
Bullet PAGE Plus Precast Gel, 15%, 17wells	R	21854-64	10 Sheets

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer
www.nacalai.com

Nucleic Acid Isolation
/ Electrophoresis

Cell Culture

Cell Extraction
/ Protein Assay

Protein Purification


Protein Electrophoresis

Western Blotting

Immunohistochemistry

DNA Ladder Markers

- » Covers wide range from 0.1 kbp to 10 kbp
- » Emphasis of 0.5, 1 and 3 kbp bands
- » Ready-to-use markers containing 2 loading dyes



Product Contents

Broad Range	100bp	1kbp
<p>DNA (ng) approx.</p> <p>10 25 8 30 6 30 5 30 4 30 3 70 2 30 1.5 35 1.2 30 1 70 0.9 30 0.8 30 0.7 30 0.6 30 0.5 90 0.4 30 0.3 40 0.2 40 0.1 50</p> <p>2% Agarose gel/TAE DNA marker 5 µl Concentration 0.15 µg/µL</p>	<p>DNA (ng) bp approx.</p> <p>1,500 30 1,200 30 1,000 70 900 30 800 30 700 30 600 30 500 90 400 30 300 40 200 40 100 50</p> <p>2.5% Agarose gel/TBE DNA marker 5 µl Concentration 0.1 µg/µL</p>	<p>DNA (ng) kbp approx.</p> <p>10 25 8 30 6 35 5 35 4 35 3 80 2 35 1.5 35 1 85 0.7 35 0.5 35 0.3 35</p> <p>0.8% Agarose gel/TAE DNA marker 5 µl Concentration 0.1 µg/µL</p>

Ordering Information

Product Name	Storage	Product No.	PKG Size
DNA Ladder One (Broad Range)(Ready-to-use)	R	08362-85	500 µl
100bp DNA Ladder One (Ready-to-use)	R	07908-75	500 µl
1kbp DNA Ladder One (Ready-to-use)	R	08232-85	500 µl

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Cell Culture

Cell Extraction
/ Protein Assay

Protein Purification

Protein Electrophoresis

Western Blotting

Immunohistochemistry







Dual Green Nucleic Acid Stain

- » Can be detected by UV excitation
- » Can be applied in both pre-stain and post-stain procedures
- » Can be used with agarose and polyacrylamide gels
- » All double-stranded DNA, single-stranded DNA and RNA can be detected.







Comparison

In agarose gel, DNA detection was performed by using post-staining and pre-staining methods. This product has equivalent or superior performance compared to products from other companies.

[Detect by UV excitation]

Staining method	Post-staining			Pre-staining		
Staining reagents	This product	Company A	Company B	This product	Company A	Company B
Staining image						

[Detect by Blue LED]

Staining method	Post-staining			Pre-staining		
Staining reagents	This product	Company A	Company B	This product	Company A	Company B
Staining image						

(Conditions)

Sample : 1kbp DNA Ladder One(Ready To Use) (#08232-85) Total nucleic acid quantity: 600 ng, 200 ng, 67 ng, 20 ng from left

Gel : 1% Agarose gel

Electrophoresis buffer : Prepare Tris-Acetate-EDTA Buffer (#35430) at 1x concentration (dye-free)

Electrophoresis : Run at 100 V until BPB reaches about 3/4 of the gel.

Staining : [Pre-staining] After dissolving the agarose, each dye was added to make a 20,000x dilution, and then the gel was solidified.

[Post-staining] TAE (1x) to make a 10,000x dilution, then place the gel after electrophoresis and shake for 30 minutes (without destaining).

Detection : ChemiDoc Touch MP(Bio-Rad laboratories) The exposure time is detected automatically.

[UV excitation] SYBR® Green mode (UV transmission, 590 / 110 nm detect on filter)

[Blue LED excitation] Alexa Fluor® 488 mode (Blue incident light, 532 / 28 nm detect on filter)

SYBR®, Alexa Fluor® is registered trademark of Molecular Probes and Incorporated.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Dual Green Nucleic Acid Stain	RT	20599-41	1 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Ethidium Bromide Solution (0.44 mg/ml)

Ethidium Bromide Solution (0.44 mg/ml) is easy and safe to use because of its eye-drop bottle. It is used in adjustment of nucleic acid staining after electrophoresis or gel containing ethidium bromide.

How to use

Adjust the concentration of ethidium bromide solution as follows

Concentration of Ethidium Bromide	Adjusting Solution	Ethidium Bromide Solution (0.44 mg/ml)
0.1 µg/ml	200 ml	1 drop
0.2 µg/ml	100 ml	1 drop
0.5 µg/ml	40 ml	1 drop



Note:

- 1 drop of Ethidium Bromide Solution (0.44 mg/ml) is 45 µl.
- In situations where Ethidium Bromide Solution (0.44 mg/ml) is used in concentrations other than shown in the above table, remove the nozzle, collect the appropriate amount with a micropipette and dilute accordingly.
- For adjustments of even greater ethidium bromide solution volumes, use Ethidium Bromide Solution (10 mg/ml) (Product No. 14631-94).

Ordering Information

Product Name	Storage	Product No.	PKG Size
Ethidium Bromide Solution (0.44 mg/ml) eye-drop-bottle	R	02393-94	10 ml
Ethidium Bromide Solution (10 mg/ml)	R	14631-94	10 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Phenol solution for DNA/RNA extraction

Ordering Information

Product Name	Details	Storage	Product No.	PKG Size
For DNA				
Phenol, Saturated with TE Buffer	Buffer-saturated phenol at pH 6.6. Easily adjusted to pH 7.9 adding the attached buffer before use.	R	25969-54 25969-96	100 ml 400 ml
Phenol, Saturated with TE Buffer	It contains 8-quinolinol as a stabilizer. The phenol layer (lower layer) is yellow in color and can be easily distinguished from the clear water layer (upper layer). pH 8 already adjusted.	R	26829-54 26829-96	100 ml 400 ml
Phenol:Chloroform:Isoamyl Alcohol 25:24:1 Mixed, pH6.7	Proteins and lipids can be removed. Easily adjusted to pH 8 adding the attached buffer before use.	R	25967-74 25967-16	100 ml 400 ml
Phenol:Chloroform:Isoamyl Alcohol 25:24:1 Mixed, pH7.9	It contains 8-quinolinol as a stabilizer. The phenol layer (lower layer) is yellow in color and can be easily distinguished from the clear water layer (upper layer). pH 8 already adjusted.	R	25970-14 25970-56	100 ml 400 ml
For RNA				
Phenol, Saturated with Citrate Buffer	Buffer-saturated phenol at pH4.3.	R	25968-64	100 ml
Phenol:Chloroform:Isoamyl Alcohol 25:24:1 Mixed, pH5.2	Proteins and lipids can be removed. Good for recovery of RNA with long poly(A).	R	26058-54 26058-96	100 ml 400 ml

Related products

Product Name	Storage	Product No.	PKG Size
Reagent for lysis			
Proteinase K from Tritirachium album	R	29442-14 29442-85	100 mg 500 mg
Proteinase K(Recombinant) Solution	R	15679-06 15679-64	2 ml 10 ml
8mol/l-Guanidine Hydrochloride Solution	RT	17356-24	100 ml
6mol/l-Guanidine Thiocyanate Solution	RT	16689-04	100 ml
100g/l-Hexadecyltrimethylammonium Bromide Solution (10% (w/v) CTAB Solution)	RT	17472-94	100 ml
Zymolyase™ 20T	R	07663-91	1 g
Zymolyase™ 100T	R	07665-55	500 mg
Buffer for electrophoresis			
Tris-Acetate-EDTA Buffer (10x) [TAE Buffer]	RT	35430-61 35430-74	1 L 5 L
Tris-Acetate-EDTA Buffer (50x) [TAE Buffer]	RT	32666-81	1 L
Tris-Borate-EDTA Buffer (5x) [TBE Buffer]	RT	35432-41	1 L
Tris-Borate-EDTA Buffer (10x) [TBE Buffer]	RT	35440-31 35440-44	1 L 5 L
Running Buffer Solution(10x) for PAGE	RT	30340-91	1 L
MOPS Buffer Stock Solution (10x) (pH 7.0)	RT	22089-45	500 ml
Reagent for polyacrilamid gel			
40(w/v)%-Acrylamide/Bis Mixed Solution(19:1), Nuclease tested	R	06140-45	500 ml
30(w/v)%-Acrylamide/Bis Mixed Solution(19:1), Nuclease tested	R	07175-75	500 ml
10(w/v)%-Ammonium Peroxodisulfate Solution	F	02634-34	10 ml
N,N,N',N'-Tetramethylethylenediamine	RT	33401-72	25 g
Denaturing reagent			
Formaldehyde Solution	RT	16223-55	500 g
Urea	RT	35940-65	500 g
Formamide	F	02020-64	100 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Nucleic Acid Isolation
/ Electrophoresis

Cell Culture

Cell Extraction
/ Protein Assay

Protein Purification

Protein Electrophoresis

Western Blotting

Immunohistochemistry

Cell Culture Medium

- » **Animal origin-free**
- » **Sterility tested for bacteria, fungus and mycoplasma**
- » **Endotoxin tested**
- » **pH tested**

Product list

Media name	Product No.	Glucose	L-Gln	Phenol Red	HEPES	Sodium Pyruvate	Other	GIBCO	Sigma
DMEM	08459	High	○	○	-	-		11965-092	D5796
	08458		○	○	-	○		11995-065	D6429
	11584		-	○	-	○		10313-021	D6546
	08457		○	○	○	-		12430-054	
	11585		-	○	○	-			D6171
	08488		-	○	-	-		11960-044	D5671
	08489		-	-	-	-		31053-028	D1145
	16971		○	○	-	○	with 1,500 mg/L sodium hydrogen carbonate		
	16972		-	○	-	-	without calcium	21068-028	
	08456	Low	○	○	-	○		11885-084	D6046
	08490		-	-	-	○		11054-020	
	09891	No	○	○	-	-		11966-025	
DMEM/ Ham's F-12	11581	○	○	○	-	○		11320-033	D8062
	08460	○	○	○	○	○		11330-032	D8437
	11583	○	-	○	○	○			D6421
	05177	○	○	-	○	○		11039-021	
	11582	○	○	-	-	○			
	09893	No	○	○	-	○			
Ham's F-12	17458	○	○	○	-	○		11765-054	N6658
G-MEM	12965	○	○	○	-	-		11710-035	
MEM	21442	○	○	○	-	-		11095-080	M4655
	21443	○	○	○	-	-	with non-essential amino acids		
	09848	No	○	○	-	-	with non-essential amino acids		
α-MEM	21445	○	○	○	-	○		12561-056	
	21444	○	○	○	-	○	with nucleosides	12571-063	
IMDM	11506	○	○	○	○	○		12440-053	I6529
RPMI 1640	30264	○	○	○	-	-		11875-093	R8758
	30263	○	○	○	○	-		22400-089	
	05176	○	-	○	-	-		21870-076	R0883
	06261	○	○	-	-	-		11835-030	R1780
	16970	○	○	○	○	○	with 4,500 mg/L glucose	A1049101	
	09892	No	○	○	-	-		11879-020	

Compositions of each product are available on online catalog, "e-Nacalai Search Version" at www.nacalai.com

For media additives, refer to p32.

Ordering Information

Product Name	Grade	Storage	Product No.	PKG size
DMEM(4.5g/l Glucose) with L-Gln, without Sodium Pyruvate, liquid	SP (For TC)	R	08459-35 08459-64	500 ml 10 x 500 ml
DMEM(4.5g/l Glucose) with L-Gln and Sodium Pyruvate, liquid	SP (For TC)	R	08458-45 08458-16	500 ml 10 x 500 ml
DMEM(4.5g/l Glucose) with Sodium Pyruvate, without L-Gln, liquid	SP (For TC)	R	11584-85	500 ml
DMEM(4.5g/l Glucose) with L-Gln and HEPES, without Sodium Pyruvate, liquid	SP (For TC)	R	08457-55	500 ml
DMEM(4.5g/l Glucose) with HEPES, without L-Gln and Sodium Pyruvate, liquid	SP (For TC)	R	11585-75	500 ml
DMEM(4.5g/l Glucose) without L-Gln and Sodium Pyruvate, liquid	SP (For TC)	R	08488-55	500 ml
DMEM(4.5g/l Glucose) without L-Gln, Sodium Pyruvate and Phenol Red, liquid	SP (For TC)	R	08489-45	500 ml
DMEM(4.5g/l Glucose) with L-Gln, Sodium Pyruvate and 1.5g/l Sodium Hydrogen Carbonate, liquid	SP (For TC)	R	16971-55	500 ml
DMEM(4.5g/l Glucose) Calcium free without L-Gln and Sodium Pyruvate, liquid	SP (For TC)	R	16972-45	500 ml
DMEM(1.0g/l Glucose) with L-Gln and Sodium Pyruvate, liquid	SP (For TC)	R	08456-65 08456-36	500 ml 10 x 500 ml
DMEM(1.0g/l Glucose) with Sodium Pyruvate, without L-Gln and Phenol Red, liquid	SP (For TC)	R	08490-05	500 ml
DMEM(No Glucose) with L-Gln, without Sodium Pyruvate, liquid	SP (For TC)	R	09891-25	500 ml
DMEM/Ham's F-12 with L-Gln and Sodium Pyruvate, without HEPES, liquid	SP (For TC)	R	11581-15	500 ml
DMEM/Ham's F-12 with L-Gln, Sodium Pyruvate and HEPES, liquid	SP (For TC)	R	08460-95	500 ml
DMEM/Ham's F-12 with Sodium Pyruvate and HEPES, without L-Gln, liquid	SP (For TC)	R	11583-95	500 ml
DMEM/Ham's F-12 with L-Gln, Sodium Pyruvate and HEPES, without Phenol Red, liquid	SP (For TC)	R	05177-15	500 ml
DMEM/Ham's F-12 with L-Gln and Sodium Pyruvate, without HEPES and Phenol Red, liquid	SP (For TC)	R	11582-05	500 ml
DMEM/Ham's F-12(No Glucose) with L-Gln and Sodium Pyruvate, liquid	SP (For TC)	R	09893-05	500 ml
Ham's F-12 with L-Gln, liquid	SP (For TC)	R	17458-65	500 ml
G-MEM with L-Gln, liquid	SP (For TC)	R	12965-65	500 ml
MEM with Earle's Salts and L-Gln, liquid	SP (For TC)	R	21442-25	500 ml
MEM with Earle's Salts, L-Gln and Non-Essential Amino Acids, liquid	SP (For TC)	R	21443-15	500 ml
MEM(No Glucose) with Earle's Salts, L-Gln and Non-Essential Amino Acids, liquid	SP (For TC)	R	09848-05	500 ml
α-MEM with L-Gln, without Ribonucleosides and Deoxyribonucleosides, liquid	SP (For TC)	R	21445-95	500 ml
α-MEM with L-Gln, Ribonucleosides and Deoxyribonucleosides, liquid	SP (For TC)	R	21444-05	500 ml
IMDM with L-Gln and HEPES, liquid	SP (For TC)	R	11506-05	500 ml
RPMI 1640 with L-Gln, liquid	SP (For TC)	R	30264-85 30264-56	500 ml 10 x 500 ml
RPMI 1640 with L-Gln and HEPES, liquid	SP (For TC)	R	30263-95	500 ml
RPMI 1640 without L-Gln, liquid	SP (For TC)	R	05176-25	500 ml
RPMI 1640 with L-Gln, without Phenol Red, liquid	SP (For TC)	R	06261-65	500 ml
RPMI 1640(4.5g/l Glucose) with L-Gln, Sodium Pyruvate and HEPES	SP (For TC)	R	16970-65	500 ml
RPMI 1640(No Glucose) with L-Gln, liquid	SP (For TC)	R	09892-15	500 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Custom Cell Culture Media

For researchers who want

- Specific compositions that appeared in the literature
- To modify the composition of commercially available cell culture media
- To get rid of phenol red due to its estrogenic effect

Specification

Product Form	Liquid
Minimum PKG Size	500 ml
Guaranteed items	pH, Osmotic pressure, Sterilized, Endotoxin tested, Mycoplasma tested
Lead terms	8-10 weeks

How to Order

Please visit our website at http://www.nacalai.co.jp/global/reagent/custom/Custom_Services.html and fill out the request form.


Reduced Serum DMEM/Ham's F-12 Medium

- » Cultivable under low-serum conditions
- » Suitable as a basal medium for serum-free cultivation
- » Reduce costs associated with serum usage
- » Minimize variations between batches, reducing serum-related effects


Comparison of cell proliferation rate and cell morphology

HEK 293 and CHO cells were cultured using both this product and Company A's serum-reduced medium. Similar to the medium from Company A, our product demonstrated that cultivation under low-serum conditions is feasible without affecting either cell proliferation rate or morphology

Proliferation curves and cell morphology of HEK 293 cells cultured under various serum concentrations (Day 35)



Proliferation curves and cell morphology of CHO cells cultured under various serum concentrations (Day 35)




Application

Cultivation of Colorectal Cancer Patient-Derived Organoids (Under Serum-Free Conditions)

We prepared droplets (n=4) embedding dispersed colon cancer patient-derived organoid cells using Corning® Matrigel Basement Membrane Matrix Growth Factor Reduced and TrypLE Express. Then, 1 ml of medium *was added on top, and the cultures were incubated.

Visual assessment of proliferation did not reveal significant differences between the two conditions.

*2 mM L-Alanyl-L-Glutamine was added to both of this product and Company A's product.



Data courtesy of Associate Professor Junpei Kondo, Department of Molecular Biochemistry, Division of Health Sciences, Graduate School of Medicine, Osaka University

Ordering Information


Product Name	Storage	Product No.	PKG Size
Reduced Serum DMEM/Ham's F-12 Medium	R	21906-55	500 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

AscleStem series

Line-up and application of AscleStem

The AscleStem® series offers a wide range of products, from undifferentiated maintenance to differentiation induction-related reagents. Using AscleStem® series products, we performed undifferentiated maintenance and differentiation induction to the trilineage in Tic line of human iPS cells.




AscleStem[®] hPSC Maintenance Supplement

AscleStem[®] hPSC Maintenance Supplement (#21979-44), is a feeder-free culture supplement for maintaining undifferentiated human pluripotent stem cells (hPSCs). It can be easily prepared by simply adding it to DMEM/Ham's F-12 medium. This product is based on the composition of a clinical grade undifferentiated maintenance medium developed through collaborative research with Osaka University. Please refer to the provided references for more information.

- » **Simply add it into DMEM/Ham's F-12 (#11581-15)**
- » **Available to reduce the cost**
- » **Applicable for the weekend skip protocol**
- » **Medium components are disclosed** * Concentration is confidential

Protocol



After thawed this product (stored in a frozen state) in a refrigerator, add the entire contents to 500 ml of DMEM/Ham's F-12. Please use it after thorough stirring.

Application


Cell morphology and alkaline phosphatase activity

The cell morphology of the 253G1 hiPSCs was confirmed after 2 weeks of culture with this product (prototype). Alkaline phosphatase staining was also performed to confirm that undifferentiated ability was not affected.

Phase contrast observation



Alkaline phosphatase staining



Reference

Hua Y, Yoshimochi K, Li J, Takekita K, Shimotsuma M, Li L, et al. Development and evaluation of a novel xeno-free culture medium for human-induced pluripotent stem cells. *Stem Cell Res Ther.* 2022;13:223. doi:10.1186/s13287-022-02879-z <https://pubmed.ncbi.nlm.nih.gov/35658933/>

Ordering Information


Product Name	Storage	Product No.	PKG Size
AscleStem [®] hPSC Maintenance Supplement	F	21979-44	32 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

AscleStem[®] Dissociation Solution for hPSCs

AscleStem[®] Dissociation Solution for hPSCs (#21777-84) is a cell dissociation solution for human pluripotent stem cells (hPSCs) composed of collagenase, trypsin, and serum substitutes. It allows for gentle dissociation of colonies during passaging of hPSCs without making single cells.


- » **Allows for gentle passaging with pipetting only**
- » **Mycoplasma tested**



<Cell detachment during passaging>
Observed passaging of human iPS cells in on-feeder culture with this product.

Application

We confirmed the undifferentiated potential of human iPS cells, which were passaged ten times using this product, by performing alkaline phosphatase staining.



201B7 on SNL76/7 feeder cells

Ordering Information


Product Name	Storage	Product No.	PKG Size
AscleStem [®] Dissociation Solution for hPSCs	F	21777-84	30 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

AscleStem® Hepatocyte Differentiation & Maintenance Basal Medium

- » Enables human pluripotent stem cells to differentiate into hepatocytes
- » Maintenance Basal Medium is suitable for drug screening applications


Protocol for hiPSC differentiation to hepatocytes



Application


Using AscleStem® Hepatocyte Differentiation/Maintenance medium, the Tic hiPSCs was induced to differentiate into hepatocytes, and albumin production was compared with that of primary cultured human hepatocytes.

Imaging: Phase microscopy



Comparison of albumin production levels

This figure shows similar albumin production levels in hepatocytes differentiated from Tic hiPSCs using AscleStem® Hepatocyte Differentiation/Maintenance medium and primary human hepatocytes cultivated 24 hours after thawing.



Ordering Information

Product Name	Storage	Product No.	PKG Size
AscleStem® Hepatocyte Differentiation Basal Medium	R	20607-05	500 ml
Supplement for H.D. Medium	F	20612-54	5 ml
AscleStem® Hepatocyte Maintenance Basal Medium	R	20683-05	500 ml
Supplement for H.M. Medium	F	20615-24	30 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

AscleStem[®] Neuronal Medium & Supplement(50x)


- » Serum-free media
- » Suitable for iPSC-derived neuronal cell culture

Differentiation from neural progenitor cells to neurons


hiPSCs-derived neural progenitor cells (day 17 after differentiation induction) were seeded at 500,000 cells/cm² and induced to differentiate into neurons by adding GlutaMAX and other inducers to our medium/our supplement or Company A's medium / their supplement. Immunostaining was performed to confirm the expression of MAP2, a marker of mature neurons.

We confirmed that hiPSC-derived neurons cultured in AscleStem[®] Neuronal Medium and AscleStem[®] Neuronal Supplement(50x) were comparable in hiPSC-derived neurons cultured in Thermo's system.

Observation by phase microscopy 26 days after differentiation




Company A's medium / their supplement




AscleStem[®] Neuronal Medium / AscleStem[®] Neuronal Supplement

Imaging by immunohistochemistry 56 days after differentiation



Company A's medium / their supplement




AscleStem[®] Neuronal Medium / AscleStem[®] Neuronal Supplement

Data courtesy of Assistant Prof. Kaneyasu Nishimura from Division of Integrated Pharmaceutical Sciences, Kyoto Pharmaceutical University


Differentiating motor neurons from hiPSCs

Using our supplement or Company A's supplement, human iPS cells were induced to differentiate into motor neurons. Immunostaining was performed to confirm the induction of motor neurons by choline acetyltransferase (ChAT) and the elongation of nerve axons by β III tubulin.


Protocol



Observed cell morphology and cell number and survival rate at detachment (28 days after start of differentiation induction)




Company A
Survival rate: 41.3%
Cell number: 4.63×10^4 cells/cm²




AscleStem® Neuronal Supplement
Survival rate: 50.2%
Cell number: 4.87×10^4 cells/cm²

IHC (β III tubulin / ChAT / DAPI : 31 days after differentiation)



Company A



AscleStem® Neuronal Supplement

Morphology, degree of axon elongation, and ChAT expression levels were comparable using AscleStem® Neuronal Supplement compared to Company A's.

Data courtesy of Stem Cell & Device Laboratory, Inc. (SCAD)

Ordering Information

Product Name	Storage	Product No.	PKG Size
AscleStem® Neuronal Medium	R	21168-35	500 ml
AscleStem® Neuronal Supplement (50x)	F	21169-54	10 ml


[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

AscleStem[®] 0.1%-Gelatin Solution


- » Ready-to-use
- » Good for cultivation of feeder cells

Application1: Adhesion of feeder cells

After coating a cell culture plate with AscleStem[®] 0.1%-Gelatin Solution, non-frozen SNL76/7 cells, cryopreserved SNL76/7 cells or MEF (CF1) cells were cultivated, then adhesion rates were measured.




Mitomycin C treated SNL76/7 cells




Application2: Maintenance culture of hiPSCs

SNL76/7 cells and MEF (CF1) cells were seeded onto plates coated with this product, on which the 253G1 hiPSCs was maintained culture and confirmed colony morphology and undifferentiated characteristics (alkaline phosphatase staining).

[Morphological observation]




253G1 on SNL76/7 feeder cells




253G1 on MEF (CF1) feeder cells

[Alkaline phosphatase staining]



253G1 on SNL76/7 feeder cells



253G1 on MEF (CF1) feeder cells

Ordering Information

Product Name	Storage	Product No.	PKG Size
AscleStem [®] 0.1%-Gelatin Solution	RT	19895-75	500 ml


[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

0.01%-Polyethyleneimine Coating Solution

- » Applicable for primary neuronal cell culture
- » Suitable for various types of cell culture labware
- » Store at room temperature

Application

Primary cultured neurons from the mouse fetal brain hippocampus were successfully maintained and cultured on various cell culture labware coated with this product.




Ordering Information

Product Name	Storage	Product No.	PKG Size
0.01%-Polyethyleneimine Coating Solution	RT	21958-14	100 ml

Mitomycin C Solution (1 mg/ml) for preparation of feeder cells

- » Ready-to-use: Sterility-tested for cell culture, does not solidify in freezer
- » High stability: 2 years in freezer, protected from light



Ordering Information

Product Name	Storage	Product No.	PKG Size
Mitomycin C Solution (1mg/ml)	F	20898-21	1 ml

Related Products

Product Name	Storage	Product No.	PKG Size
Y-27632	F	08945-71	1 mg
		08945-84	5 mg


[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Cell Dissociation Reagent: Accutase™

- » Gentle and efficient dissociation of any adherent cell line
- » No mammalian or bacterial components are contained
- » No neutralization steps by serum or trypsin inhibitors are required
- » Works extremely well on embryonic and neuronal stem cells



Application



Cell Lines dissociated with Accutase™

- hESCs
- vascular endothelial cells
- hepatocyte progenitors
- adherent CHO cells
- 293 cells
- 3T3
- HeLa
- M24 metastatic melanoma
- gliomas D54
- fibroblasts
- hepatocytes
- primary chick embryo neuronal cells
- adherent BHK cells
- L929 cells
- Vero
- NT2
- A375 metastatic melanoma
- HT1080 fibrosarcoma cells
- keratinocytes
- vascular smooth muscle cells
- bone marrow stem cells
- macrophages
- immortalized mouse testicular germ cells
- COS
- MG63
- gliomas U251
- Sf9 insect cells

Ordering Information

Product Name	Storage	Product No.	PKG Size
Accutase™	F	12679-54	100 ml

Related Products

Product Name	Storage	Product No.	PKG Size
Accumax	F	13766-74	100 ml
2.5g/l-Trypsin Solution	F	35555-54	100 ml
5.0g/l-Trypsin/5.3mmol/l-EDTA Solution	F	35556-44	100 ml
2.5g/l-Trypsin/1mmol/l-EDTA Solution	F	35554-64	100 ml
2.5g/l-Trypsin/1mmol/l-EDTA Solution, with Phenol Red	F	32777-44	100 ml
0.5g/l-Trypsin/0.53mmol/l-EDTA Solution	F	35553-74	100 ml
0.5g/l-Trypsin/0.53mmol/l-EDTA Solution, with Phenol Red	F	32778-34	100 ml
0.2g/l-EDTA Solution	R	14367-74	100 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer


EZSPHERE® for 3D cell culture with microwells

Three dimensional (3D) cell culture systems have gained in popularity as invaluable tools in broad applications of cell biology. 3D multi-cellular cell aggregates (Spheroid) can be formed by using a low attachment culture surface. However, variability in forming spheroids has been a persistent problem. EZSPHERE® is specifically designed to form a large number of uniformly sized spheroids and embryoid bodies (EBs).



- » Coated with very low binding 2-methacryloyloxyethyl phosphorylcholine (MPC) polymer
- » Form uniformly sized spheroids efficiently in the round shape wells

Well Cross-section



Ordering Information

Product Name	Storage	Product No.	PKG Size
EZSPHERE® SP, 35 mm Dish	RT	4000-900SP	10 ea
EZSPHERE® SP, 60 mm Dish	RT	4010-900SP	10 ea
EZSPHERE® SP, 100 mm Dish	RT	4020-900SP	10 ea
EZSPHERE® SP, 24-well Plate	RT	4820-900SP	5 ea
EZSPHERE® SP, 96-well Plate	RT	4860-900SP	5 ea
EZSPHERE® SP, 35 mm Dish Type 902	RT	4000-902SP	10 ea
EZSPHERE® SP, 35 mm Dish Type 903	RT	4000-903SP	10 ea
EZSPHERE® SP, 35 mm Dish Type 904	RT	4000-904SP	10 ea
EZSPHERE® SP, 35 mm Dish Type 905	RT	4000-905SP	10 ea
EZSPHERE® SP Microplate 6-well EZ-TRY	RT	TCI-4810-EZ-TRY-SP-N	5 ea
EZSPHERE® SP Microplate 24-well EZ-TRY	RT	TCI-4820-EZ-TRY-SP	5 ea

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

EZSPHERE® cell culture dishes are produced by AGC

Serum-free Cell Freezing Media: Cell Reservoir One


Cell Reservoir One is a serum-free cell culture freezing medium, which contains a water-soluble glycoprotein sericin isolated from the silkworm cocoon as a major constituent. Sericin shows the same high efficacy of cryopreservation as with FBS, and reduces the cell toxicity of DMSO. As DMSO is known to have adverse effects on cellular functions, especially embryonic stem cells, Cell Reservoir One is available both with and without DMSO.

- » No programmed freezer or special vessel necessary
- » Ready-to-use solution
- » Serum-free with no animal-derived components
- » High cell recovery and viability




Application 1

Human Mesenchymal Stem Cell (hMSC): Bone Differentiation



Application 2

Pancreatic Islet Transplantation Model



Procedure for Cell Freezing

1. Collect cells in logarithmic growth phase.
2. Suspend the cells in Cell Reservoir One (5×10^5 - 1×10^7 cells in 1 ml of Cell Reservoir One).
3. Dispense the suspension to a cryo-cell tube.
4. Store it at -80°C without pre-freezing.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Cell Reservoir One (with DMSO)	R	07485-44	100 ml
Cell Reservoir One (without DMSO)	R	07579-24	100 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

ES/iPS cell Freezing Media: Cell Reservoir One, Vitrify

Vitrification has become an important alternative to standard slow programmable freezing methods for cryopreservation of primate ES cell lines including Human iPS cells because of the higher survival rates of cells after thawing. However, the vitrification requires an ultra-rapid freezing protocol, usually less than 15 seconds between making cell suspensions and freezing in liquid nitrogen. Cell Reservoir One (Vitrify) is a novel serum-free cell culture freezing medium for vitrification method, which contains a water-soluble glycoprotein sericin isolated from the silkworm cocoon as a major constituent. It provides high survival rates of primate cells, such as Monkey ES cells and Human iPS cells even with a longer freezing protocol; up to 60 second from the cell collection to freezing in liquid nitrogen.


**Cell Reservoir One (Vitrify) is produced in corporation with SEIREN. (Patented)*

- » **High viability with a longer freezing protocol (up to 60 seconds)**
- » **Low toxicity to cells (DMSO and acetamide free)**


Application

Comparison of survival rate of Human iPS cells (201B7 cell line*) *Takahashi, K. et al. *Cell*, Nov 30;131(5):861-872 (2007)

Freezing protocol: 60 seconds



Freezing protocol: 15 seconds



Conclusion

Cell Reservoir One (Vitrify) showed high viability with both 15 and 60 seconds of freezing protocol. With 60 seconds protocol, the survival rate of cells in Cell Reservoir One (Vitrify) was significantly higher than other freezing media.

Freezing Medium		The Number of Colony		
		Vitrification Method		Slow Freezing Method
		60 Seconds	15 Seconds	
A	Cell Reservoir One (Vitrify)	672	563	-
B	DAP213	37	479	-
C	Company A	-	-	172

Ordering Information

Product Name	Storage	Product No.	PKG Size
Cell Reservoir One, Vitrify	R	11325-62	25 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Related product

Product Name	Grade	Storage	Product No.	PKG size
Dimethyl Sulfoxide	SP (For TC)	RT	13408-64	5 x 5 ml
Glycerol	SP (Biotechnology grade)	RT	09886-05	500 ml
Propylene Glycol	SP (Biotechnology grade)	RT	11740-25	500 g


Balanced Salines

» **Quality control items: pH, Osmotic pressure, Sterility test, Endotoxin test, Mycoplasma test**

D-PBS(-) w/o Ca and Mg, liquid
14249-95



D-PBS(-) w/o Ca and Mg, powder
07269-84



D-PBS(+) Preparation Reagent
(Ca, Mg Solution) (100x)
02492-94



Product list and ordering information

Product Name	KCl	Ca • Mg	Phenol Red	Storage	Product No.	PKG size	GIBCO	Sigma
D-PBS(-) without Ca and Mg, liquid	○	-	-	RT	14249-95	500 ml	14190-144	D8537
					14249-24	10 x 500 ml	14190-250	
D-PBS(-) without Ca and Mg, liquid(10x)	○	-	-	RT	11482-15	500 ml	14200-075	D1408
D-PBS without Ca and Mg, Powder	○	-	-	RT	07269-84	100 g	21600-069	D5652
Phosphate Buffered Saline without KCl(pH 7.2)	-	-	-	RT	11480-35	500 ml	20012-027	
Phosphate Buffered Saline without KCl(10x)(pH 7.2)	-	-	-	RT	11481-25	500 ml	70013-032	
Phosphate Buffered Saline without KCl(pH 7.4)	-	-	-	RT	13397-85	500 ml	10010-023	
D-PBS(+)Preparation Reagent(Ca, Mg Solution)(100x)	-	○	-	RT	02492-94	30 ml		
HBSS(+) with Ca, Mg and Phenol Red, liquid	○	○	○	RT	17459-55	500 ml	24020-117	H9269
HBSS(+) with Ca, Mg, without Phenol Red, liquid	○	○	-	RT	09735-75	500 ml	14025-092	H8264
HBSS(-) without Ca and Mg, with Phenol Red, liquid	○	-	○	RT	17460-15	500 ml	14170-112	H9394
HBSS(-) without Ca, Mg and Phenol Red, liquid	○	-	-	RT	17461-05	500 ml	14175-095	H6648

Sterile water

» **Quality control items: Sterility test, Endotoxin test, Nuclease test, Protease test**

Product list and ordering information

Product Name	Grade	Product No.	Storage	PKG size	GIBCO	Sigma
Water deionized & sterilized	SP (For molecular biology)	06442-95	RT	500 ml	15230-162	W3500
Water deionized & sterilized	SP (For molecular biology)	20620-31	F	10 x 1 ml	15230-162	-

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Media additives

● Prepared media additive solutions

» **Quality control items: Sterility test, Endotoxin test, Mycoplasma test**

Product list and ordering information

Product Name	Grade	Storage	Product No.	PKG size	GIBCO	Sigma
200mmol/l L-Alanyl-L-glutamine Solution(100x)	SP (For TC)	R	04260-64	100 ml	35050-061	G8541
200mM-L-Glutamine Stock Solution	SP (For TC)	F	16948-04	100 ml	25030-081	G7513
0.1mol/l-Calcium Chloride Solution	SP (For TC)	RT	16973-64	20 ml		
45(w/v)%-D-(+)-Glucose Solution	SP (For TC)	RT	16974-54	100 ml		G8769
1mol/l-HEPES Buffer Solution	SP (For TC)	R	17557-94	100 ml	15630-080	H0887
MEM Non-Essential Amino Acids Solution(100x)	SP (For TC)	R	06344-14 06344-56	20 ml 100 ml	11140-076 11140-050	M7145
100mM-Sodium Pyruvate Solution(100x)	SP (For TC)	R	06977-34	100 ml	11360-070	S8636

● Amino acids, vitamins and others

» **Quality control items: Endotoxin test, cell culture test**

Product list and ordering information

Product Name	Grade	Storage	Product No.	PKG size
Amino Acids				
L-α-Alanine, Animal-Free	SP (For TC)	RT	12998-32	25 g
L-Alanyl-L-glutamine, Animal-Free	SP (For TC)	R	18189-52	25 g
L-Arginine, Animal-Free	SP (For TC)	RT	11984-32 11984-45	25 g 500 g
L-(+)-Arginine Monohydrochloride, Animal-Free	SP (For TC)	RT	12999-22	25 g
L-Asparagine Monohydrate, Animal-Free	SP (For TC)	RT	13000-42	25 g
L-Aspartic Acid, Animal-Free	SP (For TC)	RT	13002-22	25 g
L-Cysteine Hydrochloride Monohydrate, Animal-Free	SP (For TC)	R	11983-42 11983-55	25 g 500 g
L-Cystine Dihydrochloride, Animal-Free	SP (For TC)	RT	13003-12	25 g
L-Glutamic Acid, Animal-Free	SP (For TC)	RT	13012-92	25 g
L-Glutamine, Animal-Free	SP (For TC)	RT	13004-02 13004-15	25 g 500 g
Glycine, Animal-Free	SP (For TC)	RT	12997-42	25 g
L-Histidine(free base), Animal-Free	SP (For TC)	RT	13014-72	25 g
L-Histidine Monohydrochloride Monohydrate, Animal-Free	SP (For TC)	RT	13017-42	25 g
L-Hydroxyproline, Animal-Free	SP (For TC)	RT	13018-32	25 g
L-(+)-Isoleucine, Animal-Free	SP (For TC)	RT	13035-02	25 g
L-Leucine, Animal-Free	SP (For TC)	RT	13036-92	25 g
L-Lysine Monohydrochloride, Animal-Free	SP (For TC)	RT	13037-82	25 g
L-Methionine, Animal-Free	SP (For TC)	RT	13038-72	25 g
L-(-)-Phenylalanine, Animal-Free	SP (For TC)	RT	13039-62	25 g
L-(-)-Proline, Animal-Free	SP (For TC)	RT	13040-22	25 g
L-Serine, Animal-Free	SP (For TC)	RT	13041-12	25 g
L-Threonine, Animal-Free	SP (For TC)	RT	13042-02	25 g
L-Tryptophan, Animal-Free	SP (For TC)	RT	13043-92	25 g
L-Tyrosine Disodium Salt Dihydrate, Animal-Free	SP (For TC)	R	11985-22 11985-35	25 g 500 g
L-Valine, Animal-Free	SP (For TC)	RT	13046-62	25 g

[Storage] RT = Room temperature, R = Refrigerator, F = Freezer

Vitamins				
p-Aminobenzoic Acid, Animal-Free	SP (For TC)	RT	13047-52	25 g
L(+)-Ascorbic Acid, Animal-Free	SP (For TC)	RT	13048-42	25 g
D-Biotin, Animal-Free	SP (For TC)	RT	13049-61	1 g
Choline Chloride, Animal-Free	SP (For TC)	R	13050-34	5 g
myo-Inositol, Animal-Free	SP (For TC)	RT	13051-82	25 g
Pyridoxine Hydrochloride, Animal-Free	SP (For TC)	RT	13053-04	5 g
Vitamin B1 Hydrochloride, Animal-Free	SP (For TC)	RT	13052-14	5 g
Vitamin B12, Animal-Free	SP (For TC)	R	13054-94	100 mg
Others				
D-(+)-Glucose, Animal-Free	SP (For TC)	RT	13057-35	500 g
Glutathione(Reduced Form), free acid, Animal-Free	SP (For TC)	R	13056-61	1 g
Hypoxanthine, Animal-Free	SP (For TC)	RT	13055-71	1 g
Sodium Pyruvate, Animal-Free	SP (For TC)	R	13058-12	25 g

● Low Molecular Compounds

(Solution)

» **Quality control items:** Sterility test, Mycoplasma test, Endotoxin test

Product list and ordering information

Product Name	Grade	Storage	Product No.	PKG size
AscleStem® CHIR99021 Solution(10mM), Animal-Free	SP (For TC)	F	21068-74	5 x 100 µL
AscleStem® SB 431542 Solution(10mM), Animal-Free	SP (For TC)	F	21235-74	5 x 100 µL
AscleStem® IWP-2 Solution(5mM), Animal-Free	SP (For TC)	F	21236-64	5 x 100 µL
AscleStem® Estradiol Solution(1mM), Animal-Free	SP (For TC)	F	21237-25	500 µL
10mmol/L-Y-27632 Solution	SP (For TC)	F	18190-96	100 µL
			18190-54	1.5 ml

(Powder)

» **Quality control items:** refer to the specification sheets for each product.

Product list and ordering information

Product Name	Grade	Storage	Product No.	PKG size
17β-Estradiol, Animal-Free	SP	R	18192-92	25 mg
	(For biochemical research)			
IWP-2, Animal-Free	SP	R	18175-64	5 mg
	(For biochemical research)			
SB 431542, Animal-Free	SP	RT	18176-54	5 mg
	(For biochemical research)			
Y-27632, Animal-Free	SP	F	18188-04	10 mg
	(For biochemical research)			

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

● Recombinant Proteins

(Solution)

» **Quality control items:** Sterility test, Endotoxin test

Product list and ordering information

Product Name	Grade	Storage	Product No.	PKG size
Activin A Solution, Human, Recombinant	SP (For TC)	F	18585-36	10 µg
			18585-94	50 µg
			18585-81	1 mg
bFGF (basic FGF) Solution, Human, Recombinant (154 a.a.), Animal-Free	SP (For TC)	F	19155-07	10 µg
			19155-36	5 x 10 µg
			19155-81	1 mg
SCF Solution, Human, Recombinant	SP (For TC)	F	21065-04	10 µg
			21065-46	50 µg
			21065-91	1 mg
KGF (FGF-7) Solution, Human, Recombinant	SP (For TC)	F	21185-34	10 µg
			21185-76	50 µg
			21185-21	1 mg
VEGF165 Solution, Human, Recombinant	SP (For TC)	F	21957-24	10 µg
			21957-66	50 µg
			21957-11	1 mg
BMP-4 Solution, Human, Recombinant	SP (For TC)	F	22247-94	10 µg
			22247-36	50 µg
			22247-81	1 mg

(Powder)

» **Quality control items:** refer to the specification sheets for each product.

Product list and ordering information

Product Name	Grade	Storage	Product No.	PKG size
Albumin, Human, Recombinant expressed in Rice, Animal-Free	SP (For TC)	F	19597-01	1 g
			19597-14	5 g
Insulin, Human, Recombinant expressed in Yeast, Animal-Free	SP (For TC)	F	12878-86	50 mg
			12878-44	250 mg
Transferrin, Human, Recombinant expressed in Rice, Animal-Free	SP (For TC)	R	12879-34	100 mg
			12879-21	1 g
			12879-76	5 g

● Proteins

(Powder)

» **Quality control items:** refer to the specification sheets for each product.

Product list and ordering information

Product Name	Grade	Storage	Product No.	PKG size
apo-Transferrin from Human	SP (For TC)	R	34401-84	100 mg
			34401-55	500 mg
Albumin, Bovine Serum, Low Endotoxin, pH5.2	SP (For biochemical research)	R	01861-84	10 g
			01861-97	100 mg
Albumin, Bovine Serum, Fatty Acid Free, pH7.0	SP (For biochemical research)	R	08587-26	10 g
			08587-42	25 g
			08587-84	50 g

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Reagents for cell dissociation

» Quality control items: pH, Sterility test, Mycoplasma test, Enzyme activity (USP unit)

Product list and ordering information

Product Name	Phenol Red	Buffer	Grade	Storage	Product No.	PKG size
0.5g/l-Trypsin/0.53mmol/l-EDTA Solution	-	HBSS(-)	SP (For TC)	F	35553-74	100 ml
0.5g/l-Trypsin/0.53mmol/l-EDTA Solution, with Phenol Red	○	HBSS(-)	SP (For TC)	F	32778-34 32778-05	100 ml 500 ml
2.5g/l-Trypsin/1mmol/l-EDTA Solution	-	HBSS(-)	SP (For TC)	F	35554-64	100 ml
2.5g/l-Trypsin/1mmol/l-EDTA Solution, with Phenol Red	○	HBSS(-)	SP (For TC)	F	32777-44 32777-15	100 ml 500 ml
5.0g/l-Trypsin/5.3mmol/l-EDTA Solution	-	HBSS(-)	SP (For TC)	F	35556-44	100 ml
2.5g/l-Trypsin Solution	-	HBSS(-)	SP (For TC)	F	35555-54	100 ml
25g/l-Trypsin Solution	-	Saline	SP (For TC)	F	18172-94	100 ml
0.2g/l-EDTA Solution	-	PBS(-)	SP (For TC)	R	14367-74	100 ml
0.5mmol/l-EDTA/PBS Solution	-	PBS(-)	SP (For TC)	RT	13567-84	100 ml
Accutase	○	PBS(-)	SP (For TC)	F	12679-54	100 ml
Accumax	-	PBS(-)	SP (For TC)	F	17087-54	100 ml

Antibiotics

Product list and ordering information

Product Name	Composition	Application	Grade	Storage	Product No.	PKG size
Actinomycin D Solution(1mg/ml)	1 mg/ml	Other	SP (For TC)	F	00393-41	1 ml
Antibiotic-Antimycotic Mixed Stock Solution(100x)	Penicillin 10,000 u/ml Streptomycin Potency: 10,000 µg/ml Amphotericin B Potency: 25 µg/ml	Bacteria, Fungi, Yeast	SP (For TC)	F	02892-54	100 ml
Antibiotic-Antimycotic Mixed Stock Solution(100x)(Stabilized)			SP (For TC)	F	09366-44	100 ml
G 418 Disulfate	-	Selection Antibiotics	SP (For TC)	RT	08973-01	1 g
					08973-14	5 g
					08973-72	25 g
G 418 Disulfate Aqueous Solution	Potency: 50 mg/ml		SP (For TC)	R	09380-86	20 ml
					09380-44	100 ml
G 418 Disulfate	-		SP (For TC)	RT	08973-01	1 g
					08973-14	5 g
G 418 Disulfate Aqueous Solution	Potency: 50 mg/ml		SP (For TC)	R	09380-86	20 ml
		09380-44			100 ml	
Gentamicin Sulfate	-	Bacteria, Mycoplasma	SP (For TC)	R	08975-81	1 g
					08975-94	5 g
Gentamicin Sulfate Solution(10mg/ml)	Potency: 10 mg/ml		SP (For TC)	R	16672-04	10 ml
Gentamicin Sulfate Solution(50mg/ml)	Potency: 50 mg/ml	SP (For TC)	R	11980-14	10 ml	
Hygromycin B	-	Selection Antibiotics	SP (For biochemical research)	R	07296-66	100 mg
					07296-11	1 g
					07296-24	5 g
Hygromycin B Solution	Potency: 50 mg/ml		SP (For TC)	R	09287-84	20 ml
Kanamycin Monosulfate	-	Bacteria, Mycoplasma	SP (For TC)	RT	08976-71	1 g
					08976-84	5 g
Kanamycin Sulfate Solution(50mg/ml)	Potency: 50 mg/ml	SP (For TC)	R	11981-04	20 ml	
Mitomycin C Solution(1mg/ml)	1 mg/ml	Other	SP (For TC)	F	20898-21	1 ml
Penicillin-Streptomycin Mixed Solution	Penicillin 10,000 u/ml Streptomycin Potency: 10,000 µg/ml	Bacteria	SP (For TC)	F	26253-84	100 ml
Penicillin-Streptomycin Mixed Solution (Stabilized)			SP (For TC)	F	09367-34	100 ml
Penicillin-Streptomycin Mixed Solution	Penicillin 5,000 u/ml Streptomycin Potency: 5,000 µg/ml		SP (For TC)	F	26252-94	100 ml
Penicillin-Streptomycin-Glutamine Mixed Solution	Penicillin 10,000 u/ml Streptomycin Potency: 10,000 µg/ml L-Glutamine 29.2 mg/ml		SP (For TC)	F	06168-34	100 ml


[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Cell Count Reagent SF, based on WST-8


- » More sensitivity than other water-soluble tetrazolium salts, such as XTT and MTS
- » No radioisotope
- » Ready-to-use




Comparison of Assay Procedure with MTT and Cell Count Reagent SF



Application for Cell Proliferation Assay



Application for Cytotoxicity Assay



References

- M. Ishiyama, Y. Miyazono, K. Sasamoto, Y. Ohkura, K. Ueno, *Talanta*, 44, 1299 (1997)
 H. Tominaga, M. Ishiyama, F. Ohseto, K. Sasamoto, T. Hamamoto, K. Suzuki and M. Watanabe, *Anal. Commun*, 36 (2), 47 (1999)

Ordering Information


Product Name	Storage	Product No.	PKG Size
Cell Count Reagent SF	R	07553-15	500 tests
		07553-44	2500 tests

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

MTT Cell Count Kit, based on reduction of MTT

- » No radioisotope
- » Ready-to-use

Cell Proliferation Assay Procedure



*1 Mix well Solubilization Solution and media as serum proteins might appear as precipitants.

*2 Avoid hard pipetting and shaking for a long time as that might help Solubilization Solution volatilize and affect the assay result.

*3 Make sure the plate is sealed completely. Alternatively, use a CO₂ incubator at 37°C.

Ordering Information


	Product Name	Storage	Product No.	PKG Size
MTT Cell Count Kit		F	23506-80	1 KIT

LDH Cytotoxicity Assay Kit

- » Economical packaging size (100 and 500-test sizes available)
- » Ready-to-use
- » Storable at 4°C for 3 months and at -20°C for 2 years without loss performance
- » By using cell suspension, enables unique assays with variety of other assays

Application

Cell damage assay with melittin



Add represented concentration of melittin to 24-hour-cultured B16/BL6 cell (1 x 10⁴ cells/well), then assay after 30-minute incubation. Substrate solution was directly added to each micro well. The higher concentration of melittin added, the stronger cell toxicity observed.

Date courtesy of Dr. Takuma Yamashita, Department of Biopharmaceutics and Drug Metabolism, Kyoto University

Ordering Information

	Product Name	Storage	Product No.	PKG Size
LDH Cytotoxicity Assay Kit		F	18250-64	100 tests
			18250-35	500 tests

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Trypan Blue Solution

» Ready-to-use



Ordering Information

Product Name	Storage	Product No.	PKG Size
Trypan Blue Solution	RT	20577-34	100 ml

Reagents for apoptosis research

Ordering Information

Product Name	Grade	Storage	Product No.	PKG size
Annexin V-633 Apoptosis Detection Kit	SP (For fluorescence analysis)	R	18146-44	100 tests
Annexin V-FITC Apoptosis Detection Kit	SP (For fluorescence analysis)	R	15342-54	100 tests
Actinomycin D Solution (1mg/ml)	SP (For TC)	F	00393-41	1 ml
Colcemid Solution (10µg/ml)	SP (For TC)	R	09356-74	10 ml
Mitomycin C Solution (1mg/ml)	SP (For TC)	F	20898-21	1 ml

Reagents for alkaline phosphatase staining

Ordering Information

Product Name	Grade	Storage	Product No.	PKG size
BCIP-NBT Solution (Ready To Use)	SP (Reagents for alkaline phosphatase staining)	R	19880-84	100 ml
4%-Paraformaldehyde Phosphate Buffer Solution	SP (For histochemistry research)	R	09154-14	5 x 10 ml
			09154-56	100 ml
			09154-85	500 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Medium for Bacteria, Plusgrow II

Plusgrow II is high performance medium for bacteria that offers easy procedures for weighing, dissolving and autoclave treatments.

- » Higher fungus density than conventional products
- » High plasmid collection

Comparison with Conventional Products

Bacteria growth test




Figure 1. Bacteria growth curve

E. coli is first cultivated in ampicillin (50 µg/ml) then added to the medium at 37°C and shaken. Then culture fluid is then centrifugally processed. Bacteria levels can then be evaluated by pellet amounts.

E. coli cell line : JM109

Plasmid : pGEM-3zf(+)



Plasmid collection test




Figure 2. Plasmid collection amount (per 1ml of culture fluid)

Plasmid is collected and purified via the alkali-SDS method (containing phenol extract) following 8 and 24 hours of *E. coli* cultivation.




Figure 3. Electrophoresis image of collection of plasmid (following 24 hours of cultivation)

Ordering Information

Product Name	Storage	Product No.	PKG Size
Plusgrow II (One package for 1L)	RT	08246-86	40 g
Dissolve → Autoclave at 121 °C for 15 minutes		08246-44	10 x 40 g
Plusgrow II	RT	08202-04	100 g
Weigh out 40 g → Dissolve in 1L → Autoclave at 121 °C for 15 minutes		08202-75	500 g

Related Products

Product Name	Storage	Product No.	PKG Size
LB Agar, Lennox	RT	20067-85	500 g
LB Agar, Miller	RT	20069-65	500 g
LB Broth, Lennox	RT	20066-95	500 g
		20066-24	2 kg
LB Broth, Miller	RT	20068-75	500 g
Agar, powder	RT	01028-85	500 g
Agar Purified, powder	RT	01162-15	500 g
Extract Yeast Dried	RT	15838-45	500 g
Tryptone	RT	35640-95	500 g

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

IPTG and X-gal Solutions

● 100mmol/l-Isopropyl-β-D-thiogalactopyranoside [IPTG] Solution


- » 0.22 μm filtrated ready-to-use solution
- » Sterilized
- » No need to adjust concentration
- » 1 ml package size allows for easy application



Application

Recombinant protein expressions are evaluated by the green fluorescent protein (GFP) expressing vector in *E. coli*.

Induced expression by IPTG



● 5-Bromo-4-chloro-3-indolyl-β-D-galactoside [X-gal] Solution (20 mg/ml)

5-Bromo-4-chloro-3-indolyl-β-D-galactoside (X-Gal) is widely used for Blue-white selection.

- » Ready-to-use DMF solution
- » 1 ml package size allows for easy application



Ordering Information

Product Name	Storage	Product No.	PKG Size
100mmol/l-Isopropyl-β-D-thiogalactopyranoside [IPTG] Solution	F	07496-91	10 x 1 ml
5-Bromo-4-chloro-3-indolyl-β-D-galactoside Solution(20 mg/ml)	F	03971-71	10 x 1 ml

Related product

Product Name	Storage	Product No.	PKG Size
Isopropyl-β-D-thiogalactopyranoside [IPTG], Dioxane free	R	19742-36	100 mg
		19742-81	1 g
		19742-94	10 g
5-Bromo-4-chloro-3-indolyl-α-D-galactoside [X-α-Gal]	R	02897-04	100 mg
		05627-86	10 mg
		05627-57	100 mg
5-Bromo-4-chloro-3-indolyl-β-D-galactoside [X-Gal]	R	05627-31	1 g
		05627-44	5 g
		05644-14	5 x 20 mg
5-Bromo-4-chloro-3-indolyl-β-D-glucuronide Cyclohexylammonium Salt	F	05646-94	10 mg
		05646-36	100 mg

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Gelling Agent for Plant Study: Gellan Gum

» High Transparency

Comparison Data with Agar Gel: Root growth observation

0.8% Agar Gel



0.2% Gellan Gum



These photos were taken on the 6th day.

Temperature : 27 °C

Light Period : 13 hours

Dark Period : 11 hours

Seeds of Komatsuna (*Brassica rapa* var. *perviridis*) were incubated in petri dishes containing a MS medium with either 0.8% agar or 0.2% Gellan Gum under sterile conditions. The seedlings were transferred to plant boxes containing the same medium 5 days later.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Gellan Gum	RT	12389-96	50 g
		12389-54	250 g

[Storage] RT = Room Temperature, _R = Refrigerator, F = Freezer

Zymolyase™ (from *Arthrobacter Luteus*)

Zymolyase™, produced by a submerged culture of *Arthrobacter luteus*⁽¹⁾, has strong lytic activity against living yeast cell walls^{(2),(3)} to produce protoplast or spheroplast of various strains of yeast cells. An essential enzyme for the lytic activity of Zymolyase™ is β -1,3-glucan laminaripentaohydrolase. It hydrolyzes linear glucose polymers with β -1,3-linkages and releases specifically laminaripentaose as the main and minimum product unit^{(4), (5), (10), (11)}. There are two preparations of Zymolyase™, Zymolyase™-20T and Zymolyase™-100T, having lytic activity of 20,000 units/g and 100,000 units/g respectively. Zymolyase™-20T is ammonium sulfate precipitate while Zymolyase™-100T is a further purified preparation by affinity chromatography⁽⁹⁾. Lytic activity varies depending on yeast strain, growth stage of yeast, or cultural conditions⁽⁶⁻⁸⁾. Further information related to Zymolyase™ can be obtained in the reference section below⁽¹²⁻¹⁶⁾.

Specifications

Product Name		Zymolyase™-20T	Zymolyase™-100T
Form		Lyophilized Powder	
Purification		Ammonium Sulfate Precipitation	Affinity Chromatography
Activity		20,000 units/g	100,000 units/g
Essential enzyme		β -1,3-glucan laminaripentaohydrolase	
Other activities contained ^(*)	β -1,3-glucanase	approx. 1.5×10^6 units/g	approx. 1.0×10^7 units/g
	protease	approx. 1.0×10^4 units/g	approx. 1.7×10^4 units/g
	mannanase	approx. 1.0×10^6 units/g	approx. 6.0×10^4 units/g
Contaminants		Amylase, Xylanase, Phosphatase	Trace amount
Optimum pH and Temp.		pH7.5, 35°C (for lysis of viable yeast cells) pH6.5, 45°C (for hydrolysis of yeast glucan)	
Stability		2°C	No loss of activity was found after storage for 1 year
Heat stability	30°C	70% of the lytic activity is lost after storage for 3 months	
	60°C	90% of the lytic activity is lost after storage for 3 months	
Specificity (Lytic Spectrum)		Lytic activity is lost on incubation for 5 minutes	
		<i>Ashbya</i> , <i>Candida</i> , <i>Debaryomyces</i> , <i>Eremothecium</i> , <i>Endomyces</i> , <i>Hansenula</i> , <i>Hanseniaspora</i> , <i>Kloeckera</i> , <i>Kluyveromyces</i> , <i>Lipomyces</i> , <i>Metschnikowia</i> , <i>Pichia</i> , <i>Pullularia</i> , <i>Torulopsis</i> , <i>Saccharomyces</i> , <i>Saccharomycopsis</i> , <i>Saccharomycodes</i> , <i>Schwanniomyces</i> , etc.	

(*)1 See reference, Kitamura, K., Kaneko, T., Yamamoto, Y., J. Gen. Appl. Microbiol., 18, 57 (1972) as to the definition of each enzyme units.

Unit Definition

One unit of lytic activity is defined as that amount which indicates 30% of decrease in absorbance at 800 nm (A_{800}) of the reaction mixture under the following condition.

[Reaction Mixture]

Enzyme solution	: 1 ml (0.05-0.1 mg/ml for Zymolyase™-20T) (0.012-0.024 mg/ml for Zymolyase™-100T)
Brewer's yeast cell suspension	: 3 ml (2 mg/ml)
1/15M Phosphate buffer	: 5 ml (pH7.5)
Distilled water	: 1 ml

After incubation for 2 hours at 25°C with gentle shaking, A_{800} of the mixture is determined. When 60% of A_{800} decrease, equivalent to 2 units, is observed in the reaction system, the brewer's yeast cells are completely lysed, namely 1 unit of Zymolyase™ lyses 3 mg dry weight of brewer's yeast.


Reference

1. Kaneko, T., Kitamura, K and Yamamoto, Y.: J. Gen. Appl. Microbiol., 15, 317 (1969)
2. Kitamura, K., Kaneko, T. and Yamamoto, Y.: Arch. Biochem. Biophys., 145, 402 (1971)
3. Kitamura, K., Kaneko, T. and Yamamoto, Y.: J. Gen. Appl. Microbiol., 18, 57 (1972)
4. Kitamura, K. and Yamamoto, Y.: Arch. Biochem. Biophys., 153, 403 (1972)
5. Kaneko, T., Kitamura, K. and Yamamoto, Y.: Agric. Biol. Chem., 37, 2295 (1973)
6. Kitamura, K., Kaneko, T. and Yamamoto, Y.: J. Gen. Appl. Microbiol., 20, 323 (1974)
7. Kitamura, K. and Yamamoto, Y.: Agric. Biol. Chem., 45, 1761 (1981)
8. Katamura, K. and Tanabe, K.: Agric. Biol. Chem., 46, 553 (1982)
9. Katamura, K.: J. Ferment. Technol., 60, 257 (1982)
10. Kitamura, K.: Agric. Biol. Chem., 46, 963 (1982)
11. Kitamura, K.: Agric. Biol. Chem., 46, 2093 (1982)
12. Calza R. E., Schroeder A. L.: J. Gen. Microbiol., 129, 413 (1983)
13. Iizuka Masaru, Torii Yasuhiko, Yamamoto Takehiko: Agric. Biol. Chem., 47 (12), 2267 (1983)
14. Shibata Nobuyuki, Kobayashi Hidemitsu, tojo Menehiro, Suzuki Shigeo: Arch. Biochem. Biophys., 251 (2), 697 (1986)
15. Iijima Y., Yanagi S. O.: Agric. Biol. Chem., 50 (7), 1855 (1986)
16. Herrero Enrique, Sanz Pascual. Sentandreu Rafael: J. Gen. Microbiol., 133 (10), 2895 (1987)

Ordering Information

Product Name	Storage	Product No.	PKG Size
Zymolyase™ 20T	R	07663-91	1 g
Zymolyase™ 100T	R	07665-55	500 mg

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer



Cell Lysis Solution: Cell Lysis Buffer

Cell Lysis Buffer is designed for protein extraction, similar to RIPA Buffer, and is available in two forms: a ready-to-use type and a 10x concentrated type. The 10x concentrated type is packaged separately with SDS solution, which may adversely affect immunoprecipitation.

- » Available for protein extraction in the same way as RIPA Buffer
- » Does not include OPE/NPE (e.g., NP-40), substances regulated under REACH
- » Available in two forms

Lineup

Available in two forms as below.


Product name	Cell Lysis Buffer	Cell Lysis Buffer (10x)
Product type	1x (ready to use)	10x
Protease inhibitor cocktail	Not included	Included
SDS	Included	Not included (10x SDS solution is packed separately)
Storage	Refrigerator	Freezer

Comparison

Protein extraction by using Cell Lysis Buffer or Company A's product

(CBB staining)

Protein was extracted from rat kidney by using this product or Company A's, then subjected to electrophoresis, and CBB staining. The result shows protein extraction efficiency of this product is equal to that of Thermo's RIPA Buffer.




<Conditions>

- Sample : ① Protein extracted by using this product (15 µg)
 : ② Protein extracted by using Company A's product (15 µg)
 : ③ Protein Ladder One Plus, Triple-color for SDS-PAGE (#19593-25)
- SDS-PAGE : Bullet PAGE Plus Precast Gel, 5-15%, 13wells (#21791-84) 400 V
- Staining : Bullet CBB Stain Lite (#21964)

(Western blotting)

Protein was extracted from mouse liver by using this product or Company A's, then subjected to western blotting. The results show that this product is applicable to western blotting, similar to Company A's.



<Conditions>

- Sample : ① Protein extracted by using this product
 : ② Protein extracted by using Company A's product
- SDS-PAGE : Bullet PAGE Plus Precast Gel, 5-15%, 13wells (#21791-84) 400 V
- Transfer : Bullet Semi-dry Transfer One (#15353-01) 25 V
- Blocking : Bullet Blocking One for Western Blotting (#13779)
- Primary antibody : [β-Actin] β-Actin antibody (C4) (Santa Cruz #sc-47778) 1,000x diluted
 [PCNA] PCNA antibody (Novus #NB100-456) 1,000x diluted
 [EGFR] EGFR antibody (CST #4267) 2,000x diluted
- Secondary antibody : [β-Actin] Anti-Mouse IgG HRP (#21860) 20,000x diluted
 [PCNA] Anti-Rabbit IgG HRP (#21858) 20,000x diluted
 [EGFR] Anti-Rabbit IgG HRP (#21858) 20,000x diluted
- Detection : Chemi-Lumi One Super (#02230)


Applications

Protein extraction by using Cell Lysis Buffer (10x)

[CBB staining]

Protein was extracted from various mouse organs by using this product with SDS, subject to electrophoresis and CBB staining.

① ② ③ ④ ⑤ ⑥ ⑦



<Conditions>

Sample : ①, ⑦ Protein Ladder One Plus, Triple-color for SDS-PAGE (#19593-25)
 ② Protein extracted from brain (12 µg)
 ③ Protein extracted from heart (12 µg)
 ④ Protein extracted from liver (12 µg)
 ⑤ Protein extracted from kidney (12 µg)
 ⑥ Protein extracted from stomach (12 µg)
 SDS-PAGE : Bullet PAGE Plus Precast Gel, 5-15%, 13wells (#21791-84) 400 V
 Stain : Bullet CBB Stain Lite (#21964)

[Immunoprecipitation]

Protein was extracted from GFP-expressing HEK293 cells using this product without SDS and immunoprecipitated using rat anti-GFP antibody-conjugated agarose beads. GFP was detected by western blotting of the samples before and after immunoprecipitation. GFP in the samples before and after immunoprecipitation was detected by western blotting.

① ②



<Conditions>

Sample : ① Sample before immunoprecipitation (3 µL)
 ② Sample after immunoprecipitation (3 µL)
 SDS-PAGE : Bullet PAGE Plus Precast Gel, 5-15%, 13wells (#21791-84) 400 V
 Transfer : Bullet Semi-dry Transfer One (#15353-01) 25 V
 Blocking : Bullet Blocking One for Western Blotting (#13779)
 Primary antibody : Anti-GFP (Mouse IgG1-κ) (#04363) 5,000x diluted
 Secondary antibody : Anti-Mouse IgG HRP (#21860) 20,000x diluted
 Detection : Chemi-Lumi One Super (#02230)

Ordering Information

Product Name	Storage	Product No.	PKG Size
Cell Lysis Buffer	R	22352-04	100 ml
Cell Lysis Buffer (10x)	F	22353-81	1 set

Related Products

Product Name	Storage	Product No.	PKG Size
Protease Inhibitor Cocktail for General Use (100x)	F	04080-11	5 x 1 ml
Protease Inhibitor Cocktail (EDTA free) (100x)	F	03969-21	5 x 1 ml
Protease Inhibitor Cocktail for Use with Mammalian Cell and Tissue Extracts	F	25955-11	5 x 1 ml
Phosphatase Inhibitor Cocktail (EDTA free)	R	07575-51	1 ml
Phosphatase Inhibitor Cocktail	R	07574-61	1 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Protease Inhibitor Cocktail

Inhibition of intra and extra cellular proteases is vital to purify and collect the expressed proteins. Saving trouble of finding adequate inhibitors, a wide range of protease is inhibited by the Protease Inhibitor Cocktail.

» **Contains inhibitors for a variety of protease**

» **Available in 3 types: General use, Mammalian cell and tissue, and EDTA free**



Composition of Each Protease Inhibitor Cocktail

Inhibitors	Target Protease	#04080-11	#03969-21	#25955-11
4-(2-Aminoethyl) benzenesulfonyl fluoride hydrochloride (AEBSF)	Serin protease	✓	✓	✓
Aprotinin	Serin protease and Esterase	✓	✓	✓
E-64	Cysteine protease	✓	✓	✓
Leupeptin hemisulfate monohydrate	Cysteine protease and Trypsin-like protease	✓	✓	✓
Disodium dihydrogen ethylenediaminetetraacetate dihydrate	Metalloprotease	✓		
Bestatin	Aminopeptidase and Leucine aminopeptidase			✓
Pepstatin A	Aspartic protease			✓

Reference

1. Okada, S. *et al. The Journal of Cellular Physiology* **226**(2), 552-558 (2011)
2. Yang, JH. *et al. The Journal of Biological Chemistry* (2010)
3. Iyama, T. *et al. Nucl. Acids Res.* **38**(14), 4834-4843 (2010)
4. Kimura, Y. *et al. Cancer Research* **70**(2), 501-511 (2010)
5. Burnett, T. J. *et al. J. Bacteriol* **165**, 139-145 (1986)
6. Hagiwara B *et al. J. Biochem.*, **45**, 185-194 (1958)

Ordering Information


Product Name	Storage	Product No.	PKG Size
Protease Inhibitor Cocktail for General Use (100x)	F	04080-11	5 x 1 ml
Protease Inhibitor Cocktail (EDTA free) (100x)	F	03969-21	5 x 1 ml
Protease Inhibitor Cocktail for Use with Mammalian Cell and Tissue Extracts	F	25955-11	5 x 1 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Phosphatase Inhibitor Cocktail

Phosphatase Inhibitor Cocktail is a mixture of several inhibitors to protect valuable proteins from dephosphorylation. The product preserves phosphorylated proteins existing in small quantity in cells and tissues.

- » Contains 6 kinds of phosphatase inhibitors for different targets
- » 100 times concentrated stock solution
- » Compatible with protein assay
- » Ready-to-use



Composition of Each Phosphatase Inhibitor Cocktail

Inhibitors	Target Phosphatase	#07575-51 EDTA free	#07574-61
Sodium orthovanadate (V)	Tyrosine phosphatase and Alkaline Phosphatase	✓	✓
Disodium molybdate (VI) dihydrate	Acid phosphatase	✓	✓
Sodium (+)-tartrate dihydrate	Acid phosphatase	✓	✓
Imidazole	Alkaline Phosphatase	✓	✓
Sodium fluoride	Acid phosphatase	✓	✓
b-Glycerophosphoric acid disodium salt	Serine-threonine phosphatase	✓	✓
tetra-Sodium ethylenediaminetetraacetate	Alkaline Phosphatase		✓

* 100 times concentrated aqueous solution

Comparison Data

Figure 1.
The detection of phosphorylated proteins in Hela cells with Anti-p-Thr antibody




Figure 2.
The phosphatase inhibition efficiency assayed by fluorescence labeled p-Tyr peptide substrate




Figure 3.
The phosphatase inhibition efficiency assayed using p-nitrophenylphosphoric acid



Reference

- Yang, JH. *et al. The Journal of Biological Chemistry* (2010)
- Selamat, W. *et al. Neuroscience Letters* **450**(2), 163-166 (2009)
- Saito, T. *et al. Biochemical and Biophysical Research Communications* **357**(2), 371-376 (2007)
- Murakami, Y. *et al. J. Biochem.*, **141**, 401-410 (2007)
- Takenaga, M. *et al. J. Cell Sci.*, **120**, 2078-2090 (2007)

Ordering Information



















Product Name	Storage	Product No.	PKG Size
Phosphatase Inhibitor Cocktail(EDTA free)	R	07575-51	1 ml
Phosphatase Inhibitor Cocktail	R	07574-61	1 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Determination of Protein Concentration: Protein Assay

The protein assay is one of the most important key techniques in proteomics. To determine protein concentration, three products with spectrophotometric method are available as follows.

Comparison of Each Method

Assay Method	Bradford	BCA	Lowry
Product Name	Protein Assay CBB Solution 	Protein Assay BCA Kit 	Protein Assay Lowry Kit 
Linearity			
	+	+++	++
Convenience	+++	++	+
Absorbance	595 nm	562 nm	750 nm
Incompatible with	Detergents	Reducing Agents	Reducing Agents
Incubate with Water			
Incubate with 0.1% SDS			
Incubate with 1 mM DTT			
Incubate with 0.1% SDS and 1 mM DTT			
Remarks	For protein samples containing detergents, BCA assay method or removal of detergents by CBB Clean Up Kit (Prod No. 11611) is helpful.	For protein samples containing reducing agents, BCA Kit with BCA Reductant Adaptable Reagent (Prod No. 21014) or the Bradford method is useful.	For protein samples containing reducing agents, the Bradford method is useful.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Protein Assay CBB Solution (Ready To Use)	R	11617-71	1 L
Protein Assay BCA Kit	RT	06385-00	1 KIT
Protein Assay Lowry Kit	RT	29470-60	1 KIT

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer


Protein Assay CBB Clean Up Kit

- » Removal interfering substances within 5 minutes
- » Designed for the Bradford protein assay



Comparison data of determination of BSA concentration with competitor's clean up kit

Protein Assay CBB Clean Up Kit is helpful to get rid of reducing agents and detergents that cause interfering with the Bradford assay, and enables better quantitative assays compared to the competitor's clean up kit.



Components

Reagent Name	Volume	Quantity
Solution A	2.5 ml	1 bottle
Solution B	2.5 ml	1 bottle
Solution C	80 ml	2 bottles
*Add 185 ml of ethanol (99.5%) into Solution C bottle, and mix it thoroughly		

Ordering Information

Product Name	Storage	Product No.	PKG Size
Protein Assay CBB Clean Up Kit	RT	11611-60	1 KIT
Protein Assay CBB Solution (Ready To Use)	R	11617-71	1 L
Albumin, Bovine Serum, Solution (2mg/ml) for Protein Assay	F	00653-31	10 x 1 ml









[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Reductant adaptable reagent for protein assay BCA Kit

- » This product enables measurement of samples containing reducing agents
- » Quick and easy: Pretreatment takes 15 mins. and measurements are completed in the same well
- » Easy dispensing and homogenizing protocols

Comparison

Measurement of samples containing reducing agents and surfactants

	PBS	0.1% SDS	1 mM DTT	0.1% SDS 1 mM DTT
BCA Kit				
Product pre-treatment operation + BCA Kit				

Effective for
samples with
both surfactants
and reductants
as well!

*Refer to the table of possible coexisting substances (instruction manual) for interfering substances that can be used in combination with reducing agents.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Protein Assay BCA Reductant Adaptable Reagent	R	21014-80	1 KIT
Protein Assay BCA Kit	RT	06385-00	1 KIT
Albumin, Bovine Serum, Solution(2mg/ml) for Protein Assay	F	00653-31	10 x 1 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

COSMOGEL[®] Ig-Accept Protein G

This product utilizes the binding of the Fc region of immunoglobulin (Ig) with Protein G, and is mainly used for the purification of IgG.

Specification

Carrier Name	COSMOGEL [®] Ig-Accept Protein G
Ligand	Protein G
Particle size	50 - 150 μm
Matrix	4% Cross-linked agarose
Recommended Line Velocity	26 cm/h
Limit pressure	3.6 psi (0.25 bar)
Supply condition	50 vol% suspension (20 vol% ethanol solution)

Application

COSMOGEL[®] Ig-Accept Protein G



Sample : Mouse serum (ammonium sulfate precipitation fraction)
Elution : Glycine - Hydrochloric acid buffer (pH 2.8)
Lane : 1. Protein Markers (10x) (#29458-24) in 1x
2. Sample purified by COSMOGEL[®] Ig-Accept Protein G

Ordering information

Product Name	Storage	Product No.	PKG Size
COSMOGEL [®] Ig-Accept Protein G	R	02198-64	5 ml
		02198-22	25 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Nucleic Acid Isolation
/ Electrophoresis

Cell Culture

Cell Extraction
/ Protein Assay

Protein Purification

Protein Electrophoresis

Western Blotting

Immunohistochemistry


COSMOGEL® GST-Accept

- » Superior durability
- » Superior cost performance

Application

10 consecutive purifications of GST-tag fusion proteins

We performed 10 consecutive purifications of GST-His fusion proteins by using the same resin from *E. coli* lysate. The 10th purification was also successful.



Sample : *E. coli* (BL21(DE3)pLysS) lysate transformed with GST-His(pET-41b(+))
 Binding buffer : Use Phosphate Buffered Saline(10x)(pH 7.4)(#27575-31)1x
 Elution buffer : 10 mM Glutathione, 50 mM Tris-HCl, pH 8.0

Ordering information

Product Name	Storage	Product No.	PKG Size
COSMOGEL® GST-Accept	R	09277-56	5 ml
		09277-72	25 ml
		09277-14	100 ml


[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

High Performance Magnetic Nanoparticles: FG beads®


- » Excellent recovery of target proteins
- » Extremely low non-specific binding
- » High stability in organic solvents

The FG beads developed by Tokyo Institute of Technology consists of ferrite nanoparticles coated firmly with a polymer layer and its diameter is approx. 200 nm. The FG beads are used as carriers for affinity purification of target proteins.¹⁾

Purification Process



Comparison with Other Magnetic Beads²⁾



1. S. Sakamoto et al., *Chem. Rec.* 9 (2009) 66

2. K. Nishio et al., *Colloids Surfaces. B.* 64 (2008) 162

Ordering Information

Product Name	Density	Storage	Product No.	PKG Size
Plain beads	20mg/ml	R	TAS8848N1010	0.5ml
Linker beads (Epoxy beads)	20mg/ml	R	TAS8848N1110	0.25ml
OH beads	20mg/ml	R	TAS8848N1120	0.25ml
NH2 beads	20mg/ml	R	TAS8848N1130	0.25ml
COOH beads	20mg/ml	R	TAS8848N1140	0.25ml
NHS beads	20mg/ml	F	TAS8848N1141	0.25ml
Ts beads	20mg/ml	R	TAS8848N1150	0.25ml
Azide beads	20mg/ml	R	TAS8848N1160	0.25ml
Alkyne beads	20mg/ml	R	TAS8848N1161	0.25ml
Streptavidin beads	20mg/ml	R	TAS8848N1170	0.25ml
NeutrAvidin beads	20mg/ml	R	TAS8848N1171	0.25ml
Protein A beads	20mg/ml	R	TAS8848N1172	0.25ml
Protein G beads	20mg/ml	R	TAS8848N1173	0.25ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

FG beads® are produced by Tamagawa Seiki Co., Ltd.



Product Name	Density	Storage	Product No.	PKG Size
HM-Streptavidin beads	20mg/ml	R	TAB8848N3170	0.25ml
HM-NeutrAvidin beads	20mg/ml	R	TAB8848N3171	0.25ml
HM-Protein A beads	20mg/ml	R	TAB8848N3172	0.25ml
HM-Protein G beads	20mg/ml	R	TAB8848N3173	0.25ml
Magnetic Stand (for 1.5 ml tube)		RT	TAB4899N12	1ea
Magnetic Stand (for 15 ml tube)		RT	TAB4899N20	1ea
Magnetic Stand (for 50 ml tube)		RT	TAB4899N30	1ea
Magnetic Stand (for PCR tube)		RT	TAB4899N41	1ea

FG beads® are produced by Tamagawa Seiki Co., Ltd.

TurboTEV Protease & Turbo3C Protease

- » Both GST and His tags to facilitate its removal from the digested protein sample
- » Activity over a broad temperature (4°C to 37°C) and pH (6.5 to 8.5) range

Application



GST-fusion protein (C) at 1 mg/ml is incubated with TurboTEV or Turbo3C Protease at a ratio of (1) 1:50, (2) 1:100, (3) 1:200 (w/w) in a buffer of 25 mM Tris-HCl, pH 8.0, 150 mM NaCl, 14 mM 2-mercaptoethanol at 4°C for 16 hours.

Specification

	TurboTEV Protease	Turbo3C Protease
Molecular Weight	52 kDa	47 kDa
Cleavage Site	Glu-Asn-Leu-Tyr-Phe-Gln↓Gly	Leu-Glu-Val-Leu-Phe-Gln↓Gly-Pro
Enzymatic Activity	One unit cleaves >85% of 3 µg control substrate at 30°C for 1 h	One unit cleaves >95% of 100 µg control substrate at 4°C for 16 h
Concentration	20,000 units/ml	2,000 units/ml
Cleavage Condition	A broad temperature (4°C to 37°C) and pH (6.5 to 8.5) range	
Formulation	25 mM Tris-HCl(pH8.0), 50 mM NaCl, 1 mM TCEP, 50% Glycerol	

Ordering Information

Product Name	Storage	Product No.	PKG Size
TurboTEV (TEV Protease) 2 mg/ml	F	NU0102S	1,000 units (0.1 mg)
		NU0102M	10,000 units (1 mg)
		NU0102L	100,000 units (10 mg)
Turbo3C (HRV3C Protease) 2 mg/ml	F	NU0101S	1,000 units (1 mg)
		NU0101M	10,000 units (10 mg)

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

TurboTEV Protease & Turbo3C Protease are produced by Nacalai USA.

[Ultra-fast Precast Gel]




Bullet PAGE Plus Precast Gel

- » Only 10 minutes with 400 V
- » Wide range of gel concentrations available
- » Compatible with conventional Laemmli running buffer and sample buffer
- » Surfactant-free, so can be used for nucleic acid analysis (refer to p11)

*Please use 25 mM Tris and 192 mM glycine buffer for nucleic acid electrophoresis

Performance Comparison

This product demonstrates separation ability and band sharpness comparable to competitors' gels, despite a short running time of about 10 minutes.

Product Name	Bullet PAGE Plus	Company A	Company B
Gel %	5 - 20%	4 - 20 %	4 - 12%
Prod No.	21794-54	-	-
Running Time	10 m 07 s	29 m 35 s	48 m 32 s
Separation Image			

<Condition>

Sample : Protein Markers(10x)(# 29458-24), 3μl


Gel Staining : CBB Stain One (#04543)

Voltage Constant : Bullet PAGE Plus: 400V






Company A and Company B: 200V





Blotting performance: Comparison of protein transfer efficiency

This product shows a transfer efficiency equivalent to Company A's product.



Gel types

Gel Type	Gradient Gel				
Gel %	5 - 10%	5 - 15%	5 - 20 %	7.5 - 15 %	10 - 20 %
Prod No. for 13-well for 17-well	21789-34 21790-94	21791-84 21792-74	21793-64 21794-54	21795-44 21796-34	21797-24 21798-14
Separation Image					

Gel Type	Single-percentage gel			
Gel %	7.5%	10%	12.5%	15%
Prod No. for 13-well for 17-well	21799-04 21800-54	21801-44 21806-94	21807-84 21811-14	21853-74 21854-64
Separation Image				

Specification

Glass Plate Size

: W100 mm x H80 mm x T3.2 mm (made of glass)

Gel Size

: W80 mm x H60 mm x T1.0 mm

Sample Well Configuration / Maximum Load Volume

: 13-well / 40 μ l, 17-well / 28 μ l

Shelf Life

: 9 months

Nucleic Acid Isolation / Electrophoresis
Cell Culture
Cell Extraction / Protein Assay
Protein Purification
Protein Electrophoresis
Western Blotting
Immunohistochemistry

Ordering Information

Product name	Storage	Product number	PKG size
Bullet PAGE Plus Precast Gel, 5-10%, 13wells	R	21789-34	10 Sheets
Bullet PAGE Plus Precast Gel, 5-10%, 17wells	R	21790-94	10 Sheets
Bullet PAGE Plus Precast Gel, 5-15%, 13wells	R	21791-84	10 Sheets
Bullet PAGE Plus Precast Gel, 5-15%, 17wells	R	21792-74	10 Sheets
Bullet PAGE Plus Precast Gel, 5-20%, 13wells	R	21793-64	10 Sheets
Bullet PAGE Plus Precast Gel, 5-20%, 17wells	R	21794-54	10 Sheets
Bullet PAGE Plus Precast Gel, 7.5-15%, 13wells	R	21795-44	10 Sheets
Bullet PAGE Plus Precast Gel, 7.5-15%, 17wells	R	21796-34	10 Sheets
Bullet PAGE Plus Precast Gel, 10-20%, 13wells	R	21797-24	10 Sheets
Bullet PAGE Plus Precast Gel, 10-20%, 17wells	R	21798-14	10 Sheets
Bullet PAGE Plus Precast Gel, 7.5%, 13wells	R	21799-04	10 Sheets
Bullet PAGE Plus Precast Gel, 7.5%, 17wells	R	21800-54	10 Sheets
Bullet PAGE Plus Precast Gel, 10%, 13wells	R	21801-44	10 Sheets
Bullet PAGE Plus Precast Gel, 10%, 17wells	R	21806-94	10 Sheets
Bullet PAGE Plus Precast Gel, 12.5%, 13wells	R	21807-84	10 Sheets
Bullet PAGE Plus Precast Gel, 12.5%, 17wells	R	21811-14	10 Sheets
Bullet PAGE Plus Precast Gel, 15%, 13wells	R	21853-74	10 Sheets
Bullet PAGE Plus Precast Gel, 15%, 17wells	R	21854-64	10 Sheets

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer







[General-use Precast Gel]

Extra PAGE One Precast Gel

- » **Superior resolution**
- » **Sharp bands**
- » **High transfer efficiency**
- » **Compatible with conventional Laemmli sample and running buffers**

Performance Comparison

Extra PAGE One Precast Gel offers higher resolution than other gels. Especially 5-20% gradient gel and 12.5% gel provide improved resolution for low molecular weight.

Gel	Gradient Gel				Single-percentage Gel	
Product Name	Extra PAGE One	Company A	Extra PAGE One	Company B	Extra PAGE One	Company A
Gel%	5 - 20%	4 - 20%	5 - 20%	5 - 20%	12.5%	12.5%
Product No.	11955-54	-	11955-54	-	11965-24	-
Migration patterns						
	M.W.(KDa) 200→ 45→ 22→ 6.5→	M.W.(KDa) 200→ 45→ 22→ 6.5→	M.W.(KDa) 200→ 45→ 22→ 6.5→	M.W.(KDa) 200→ 45→ 22→ 6.5→	M.W.(KDa) 200→ 45→ 22→ 14→ 6.5→	M.W.(KDa) 200→ 45→ 22→ 6.5+14→

<Condition>


Sample : Protein Markers (#29458-24) 5ul.

Staining : CBB Stain One (#04543)

Electrophoresis condition : Extra PAGE One Precast Gel: 300 V, approx. 50 min.
Company A: followed the manufacture's instructions

Comparison of Transfer Efficiency

Extra PAGE One Precast Gel offers higher transfer efficiency than Company A for both high and low molecular weight proteins.



<Sample>

1. Protein ladder One (Triple-color) (Product No. 09547)
2. 2 ul of Protein Markers (3x)
3. 2 ul of Protein Markers (1x)
4. 2 ul of Protein Markers (1/3)
5. 10 ul of Bio-Rad Precision Plus Protein™, Kaleidoscope™

<Transfer>






Semi-dry with PVDF membrane and Blotting Buffer Solution for Western Blotting (Product No.: 30650-31) (Transfer time 20 min with 10V constant current)





<Staining>

CBB Stain One (Product No.: 04543)

Product Lines

Extra PAGE One Precast Gel is available in gradient gels and single-percentage gels.

Gel	Gradient Gel				
Gel %	5 - 10%	5 - 15%	5 - 20%	7.5 - 15 %	10 - 20 %
Product No. 13well/17well	13059-44 / 13060-04	13061-94 / 13062-84	13063-74 / 13064-64	13065-54 / 13066-44	13067-34 / 13068-24
Migration patterns	M.W.(kDa) 	M.W.(kDa) 	M.W.(kDa) 	M.W.(kDa) 	M.W.(kDa) 
	200→ 45→	200→ 45→ 22→ 6.5→	200→ 45→ 22→ 6.5→	200→ 45→ 22→ 6.5→	200→ 45→ 22→ 6.5→

Gel	Single-percentage Gel			
Gel %	7.5%	10%	12.5%	15%
Product No. 13well/17well	13069-14 / 13070-74	13071-64 / 13072-54	13073-44 / 13074-34	13075-24 / 13076-14
Migration patterns	M.W.(kDa) 	M.W.(kDa) 	M.W.(kDa) 	M.W.(kDa) 
	200→ 45→	200→ 45→ 22→	200→ 45→ 22→ 6.5→	200→ 45→ 22→ 6.5→

Note) Back ground may remain slightly when using Nacalai's Rapid Stain CBB Kit (Product No.: 30035-14) or other products. To reduce back ground, wash the gel with 10% acetic acid for 20 minutes.

Specifications

Cassette Size : 100 mm(W) x 100 mm(H) x 3.2 mm(T) (made of glass)

Gel Size : 80 mm(W) x 80 mm(H) x 1.0 mm(T)

Well : 13 wells, 17 wells

Shelf Life Expire date : 9 months

Note) Use electrophoresis tank for plate size 100mm(W) x 100mm(H) x 3.2mm(T).

Ordering Information

Product name	Storage	Product number	PKG size
Extra PAGE One Precast Gel, 5-10%, 13wells	R	13059-44	10 Sheets
Extra PAGE One Precast Gel, 5-10%, 17wells	R	13060-04	10 Sheets
Extra PAGE One Precast Gel, 5-15%, 13wells	R	13061-94	10 Sheets
Extra PAGE One Precast Gel, 5-15%, 17wells	R	13062-84	10 Sheets
Extra PAGE One Precast Gel, 5-20%, 13wells	R	13063-74	10 Sheets
Extra PAGE One Precast Gel, 5-20%, 17wells	R	13064-64	10 Sheets
Extra PAGE One Precast Gel, 7.5-15%, 13wells	R	13065-54	10 Sheets
Extra PAGE One Precast Gel, 7.5-15%, 17wells	R	13066-44	10 Sheets
Extra PAGE One Precast Gel, 10-20%, 13wells	R	13067-34	10 Sheets
Extra PAGE One Precast Gel, 10-20%, 17wells	R	13068-24	10 Sheets
Extra PAGE One Precast Gel, 7.5%, 13wells	R	13069-14	10 Sheets
Extra PAGE One Precast Gel, 7.5%, 17wells	R	13070-74	10 Sheets
Extra PAGE One Precast Gel, 10%, 13wells	R	13071-64	10 Sheets
Extra PAGE One Precast Gel, 10%, 17wells	R	13072-54	10 Sheets
Extra PAGE One Precast Gel, 12.5%, 13wells	R	13073-44	10 Sheets
Extra PAGE One Precast Gel, 12.5%, 17wells	R	13074-34	10 Sheets
Extra PAGE One Precast Gel, 15%, 13wells	R	13075-24	10 Sheets
Extra PAGE One Precast Gel, 15%, 17wells	R	13076-14	10 Sheets

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer


[Apparatus for Precast Gel]

Electrophoresis Tank for Precast Gel

Specification

Size : 143W x 84D x 140H (mm)

Required Buffer Volume : 550 ml



Ordering Information

Product Name	Storage	Product No.	PKG Size
WEP-MN Vertical Electrophoresis Tank	RT	WEP-MN	1 Set

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Note: Since the adapter for fitting Bullet PAGE Plus includes this product, please inform a distributor if you plan to use this product for Extra PAGE One.

[Gel Buffer for Hand-made Gel]

WIDE RANGE Gel Preparation Buffer (4x) for PAGE

Gradient gels offer a much wider separation range of proteins than single percentage gels. However, casting gradient gels is more difficult and labor intensive. WIDE RANGE Gel Preparation Buffer offers a gradient gel-like separation on a single percentage gel by simply mixing it with acrylamide/ bisacrylamide gel casting solution. The gel can be used with the common sample buffers and running buffers. It is also suitable for standard staining methods including CBB and silver staining.


» Simple casting procedure

WIDE RANGE Gel Preparation buffer is a 4x concentrated neutral pH buffer. It can be used for preparation of both stacking gel and separation gel by replacing the Tris-HCl buffer in Laemmli buffer system.




» Improved stability and strength

The increased tensile strength allows easy handling even a low percentage gel. The neutral pH buffer improves the stability of gel resulting in a longer shelf life than the gel with Laemmli buffer system.




» A wide separation range

WIDE RANGE Gel provides a much greater separation range than the gel casted with a conventional Laemmli buffer system.



Applicable for Western Blotting




Ordering Information

Product Name	Storage	Product No.	PKG Size
WIDE RANGE Gel Preparation Buffer (4x) for PAGE	R	07831-94	250 ml

[Stacking Gel Buffer for Hand-made Gel]

Stacking Gel Buffer Solution (4x) with Blue Color

» Easy to see well locations due to coloring stacking gel



Wells are clearly confirmed on stacking gel prepared with Stacking Gel Buffer Solution with Dye.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Stacking Gel Buffer Solution (4x) with Dye for SDS-PAGE	R	09268-34	100 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

[Fast running buffer for Hand-made Gel]



Rapid Running Buffer Solution

- » Approximately 25 min. electrophoresis time with mini-gel at 250 V
- » Just replace the Laemmli running buffer with this product
- » High protein transfer efficiency to western blotting membrane



Comparison of electrophoresis duration and buffer temperature rise

Running proteins with this product shortens the electrophoresis duration to about 60% compared to Laemmli running buffer, and its rising in buffer temperature is the almost same as Laemmli running buffer's.



Separation patterns of Laemmli gel with Rapid Running Buffer

Gel concentration	6%	8%	10%	4-10%
M.W.(kDa)	200→ 45→ 22→	200→ 45→ 22→ 6.5→	200→ 45→ 22→ 6.5→	200→ 45→ 22→ 6.5→
Separation image				

Comparison of protein transfer efficiency to western blotting membrane

Due to lower gel concentration when Rapid Running Buffer Solution is used^{*1}, its protein transfer efficiency to a membrane is higher than the Laemmli running buffer's.

^{*1} By casting a gel with 4% lower gel concentration than usual, its separation patterns can be made similar to the original's, e.g. separation patterns of 8% gel with Rapid Running Buffer Solution is about the same as 12% gel with Laemmli running buffer.

Gel concentration	8%	12%	8%	12%
Running buffer	Rapid Running Buffer	Laemmli running buffer	Rapid Running Buffer	Laemmli running buffer
Blotting time	10 min.			
M.W.(kDa)	1 2 3 4	1 2 3 4	1 2 3 4	1 2 3 4
200→				
45→				
22→				
6.5→				
Membrane staining image				

Ordering Information

Product Name	Storage	Product No.	PKG Size
Rapid Running Buffer Solution (20x) for SDS-PAGE	RT	12981-74	250 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer



Polyacrylamide Gel Casting Reagents

Ordering Information

Product Name	Storage	Product No.	PKG Size
Acrylamides (monomer)			
Acrylamide (monomer), Purity, 99%	RT	00809-14	100 g
		00809-85	500 g
Acrylamide (monomer), Purity, 99%, Nuclease and Protease tested	RT	06114-24	100 g
		06114-95	500 g
		06114-11	1 kg
Crosslinking Agents			
N,N'-Methylenebisacrylamide, [BIS]	R	22402-02	25 g
N,N'-Methylenebisacrylamide, Purity, 99%, Nuclease and Protease tested	R	22407-52	25 g
Acrylamide/Bis Mixed Solutions			
30(w/v)%-Acrylamide/Bis Mixed Solution (37.5:1)	R	06144-05	500 ml
30(w/v)%-Acrylamide/Bis Mixed Solution (29:1)	R	06141-35	500 ml
40(w/v)%-Acrylamide/Bis Mixed Solution (37.5:1)	R	06121-95	500 ml
40(w/v)%-Acrylamide/Bis Mixed Solution (29:1)	R	06119-45	500 ml
Polymerization Initiators			
N,N,N',N'-Tetramethylethylenediamine [TEMED]	RT	33401-72	25 g
		33401-14	100 g
Polymerization Promotors			
Ammonium Peroxodisulfate [APS]	R	02627-21	1 g
		02627-34	10 g
10 (w/v)%-Ammonium Peroxodisulfate Solution	F	02634-34	10 ml
Gel Buffer Solutions			
Separating Gel Buffer Solution (4x) for SDS-PAGE, pH8.8 Filtrated by 0.45 µm filter Components : 1.5M-Tris-HCl, 0.4 (w/v)%-SDS	RT	30651-05	500 ml
Stacking Gel Buffer Solution (4x) with Dye for SDS-PAGE, pH6.8 Filtrated by 0.45 µm filter Components : 0.5M-Tris-HCl, 0.4(w/v)%-SDS, Dye	R	09268-34	100 ml
Stacking Gel Buffer Solution (4x) for SDS-PAGE, pH6.8 Filtrated by 0.45 µm filter Components : 0.5M-Tris-HCl, 0.4 (w/v)%-SDS	R	09267-44	100 ml
		32158-25	500 ml

Running Buffers

Ordering Information

Product Name	Storage	Product No.	PKG Size
Pre-mixed Buffers			
Running Buffer Solution (10x) for SDS-PAGE,Tris-Glycine, Filtrated by 0.45 µm filter Components: 0.25 mol/l-Tris, 1.92 mol/l-glycine, 10 g/l-SDS	RT	30329-61 30329-74	1 L 5 L
Running Buffer Solution (10x) for PAGE,Tris-Glycine, Filtrated by 0.45 µm filter Components: 0.25 mol/l-Tris, 1.92 mol/l-glycine	RT	30340-91	1 L
Buffer Adjusting Reagents			
Tris(hydroxymethyl)aminomethane, Purity, 99%	RT	35410-34	100 g
Tris(hydroxymethyl)aminomethane, Purity, 99.9%, Nuclease and Protease tested	RT	35434-76	100 g
		35434-05	500 g
		35434-21	1 kg
Glycine	RT	17128-14	100 g
Glycine, Nuclease and Protease tested	RT	17141-24	100 g
		17141-95	500 g
Tricine {N-[Tris(hydroxymethyl)methyl]glycine}	RT	34713-62	25 g
		34713-04	100 g
Tricine {N-[Tris(hydroxymethyl)methyl]glycine} Nuclease and Protease tested	RT	02437-24	100 g
Sodium Lauryl Sulfate [Sodium Dodecyl Sulfate;SDS] Purity, 99%	RT	31607-52	25 g
		31607-94	100 g
		31607-65	500 g
Sodium Lauryl Sulfate granular [Sodium Dodecyl Sulfate;SDS] Purity, 99%, Solids (granular)			
<div></div>	RT	02873-62	25 g
		02873-04	100 g
		02873-75	500 g
10%-SDS Solution [10%-Sodium Lauryl Sulfate Solution]	RT	30562-04	100 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Sample Buffer Solution for SDS-PAGE (6x)

- » Suitable for low concentration protein sample adjustment
- » No precipitation in the refrigerator
- » Two types of reagents (with and without reducing agent)

Ordering Information

Product Name	Storage	Product No.	PKG Size
Sample Buffers			
Sample Buffer Solution with Reducing Reagent (6x) for SDS-PAGE pH6.8 Filtrated by 0.45 µm filter, Components: 0.375M-Tris-HCl, 0.03(w/v)%-BPB, glycerin, anion surface acting agent and reducing agent	R	09499-14	5 ml
Sample Buffer Solution without Reducing Reagent (6x) for SDS-PAGE pH6.8 Filtrated by 0.45 µm filter, Components: 0.375M-Tris-HCl, 0.03(w/v)%-BPB, glycerin and anion surface acting agent	R	09500-64	5 ml
Sample Buffer Solution with 2-ME (2x) for SDS-PAGE pH6.8 Filtrated by 0.45 µm filter, Components: 0.125M-Tris-HCl, 4(w/v)%-SDS, 20(v/v)%-glycerin, 0.01(w/v)%-BPB, 10(v/v)%-2-ME	R	30566-22	25 ml
Sample Buffer Solution without 2-ME (2x) for SDS-PAGE pH6.8 Filtrated by 0.45 µm filter, Components: 0.125M-Tris-HCl, 4(w/v)%-SDS, 20(v/v)%-glycerin, 0.01(w/v)%-BPB	R	30567-12	25 ml
Reducing Agent			
2-Mercaptoethanol	RT	21418-42	25 g
		21418-84	100 g
		21418-55	500 g
Dithiothreitol	R	14112-36	100 mg
		14112-81	1 g
		14112-94	5 g
		14112-52	25 g
1mol/l-Dithiothreitol Solution	F	14130-41	1 ml
Tris (2-carboxyethyl) phosphine Hydrochloride (TCEP)	R	07277-61	1 g
Tracking Dyes			
Bromophenol Blue	RT	05808-61	1 g
		05808-32	25 g
Others			
Glycerol Nuclease and Protease tested	RT	17045-94	100 ml
		17045-65	500 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Nucleic Acid Isolation
/ Electrophoresis

Cell Culture

Cell Extraction
/ Protein Assay

Protein Purification

Protein Electrophoresis


Western Blotting

Immunohistochemistry

Molecular Weight Markers

● Protein Ladder One Plus

- » Excellent visibility with triple color
- » Sharp bands
- » Consists of 14 bands
- » Low lot-to-lot variation



Extra PAGE One Precast Gel, 5-20%, 13wells(#13063-74)
Load volume 5 μ L

Notes


- There is a possibility of different molecular weight result depending on gel type (concentration, gradient, buffer, etc.). It may not be suitable for accurate molecular weight measurement.
- The product is ready-to-use type. Sample preparation, such as heating and reduction, is not required.

Protein	
14 kinds	
Composition	
Glycerol	15% (v/v)
SDS	2% (w/v)
DTT	0.2 mM
Urea	3.6 M
Tris-H ₃ PO ₄	20 mM
pH 7.5	

● Unstained Protein Markers (10x)

- » Contains 8 kinds of protein (M.W. 6,500 - 200,000 Da)

Image



Composition


50(v/v)% Glycerol
0.3 M NaCl
0.1 M DTT, 2 mM EDTA \cdot 2Na
3 mM NaN₃
10 mM Tris-HCl (pH 7.0)

●Chemi-Lumi One Markers Kit


Chemi-Lumi One Markers Kit consists of biotinylated proteins and HRP-conjugated streptavidin. Each band can be visualized on a western blotting by the same chemiluminescent reagents for the target protein.

» **Contains 8 biotinylated proteins as molecular weight markers (M.W. 6,500 - 200,000 Da)**

» **Includes HRP-conjugated streptavidin to detect biotinylated proteins**



Western Blotting for Detecting of β -Actin



Components

Chemi-Lumi One Markers consists of 8 biotinylated proteins, 50 μ l: 1 tube

Streptavidin (HRP conjugate), 250 μ l: 1 tube

Note: The molecular weight of Chemi-Lumi One Markers may slightly differ from unmodified proteins because of biotinylation.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Protein Ladder One Plus, Triple-color for SDS-PAGE	F	19593-25	500 μ L
Protein Markers(M.W. 6,500~200,000)(10x) for SDS-PAGE	F	29458-24	200 μ L
Chemi-Lumi One Markers Kit	F	06456-70	1 KIT


[Storage] RT = Room temperature, R = Refrigerator, F = Freezer

PAGE Clean Up Kit

This product is designed for sample preparation in two-dimensional electrophoresis and SDS-PAGE. This product is a combination of acetone, trichloroacetic acid, and coprecipitate optimized for protein precipitation. By resuspending the precipitated proteins in sample lysate according to the intended use, it is possible to concentrate the sample protein and remove interfering substances in the buffer.

- » **Suitable for concentrating protein samples and removing analytical interferences**
- » **Precipitation is possible even in the subsistence of high concentrations of urea, guanidine, and SDS**
- » **Easier than dialysis and small volume processing at the μ l level**

Removing electrolyte interfering substances



<Conditions>

[Sample]

- (A) 10x dilution of Protein Markers(10x) #29458-24 (standard concentration)
- (B) BSA (100ng: non-reducing)

[Precipitation treatment]

- ① without this product
- ② with this product

[Electrophoresis]

SDS-PAGE (12% gel)

[Staining]

- (A) CBB Stain One #04543 (CBB staining)
- (B) Sil-Best Stain One #06865-81 (silver staining)

When a sample containing components that affect the electrophoretic image was treated with this product, the electrophoretic image was improved.

Note) Yield may vary depending on the composition of the sample solution, protein concentration, and other factors. Please use after preliminary examination.

Ordering Information

Product Name	Storage	Product No.	PKG Size
PAGE Clean Up Kit	R	06441-50	1 KIT

[Storage] RT = Room temperature, R = Refrigerator, F = Freezer

CBB Stain Solution

CBB staining, Coomassie Brilliant Blue staining, is a popular method for detecting proteins in polyacrylamide gel. We offer four types of CBB staining solution.

Selection of CBB staining solution

	Bullet CBB Stain Lite (#21964)	CBB Stain One Super (#11642)	CBB Stain One (#04543)	Rapid Stain CBB Kit (#30035)
Dye Type	CBB-G250	CBB-R250	CBB-G250	CBB-R250
Component	Single bottle (Ready-to-use) w/o acetic acid and methanol	Single bottle (Ready-to-use) w/o acetic acid and methanol	Single bottle (Ready-to-use) w/o acetic acid and methanol	Two bottles w/ acetic acid and w/o methanol
Gel rinsing	Unnecessary	Required 3 times for 5min.	Required 3 times for 5min.	Unnecessary
Staining Period	15 min.	30 min.	60 min.	20 min.
Destaining Period	Unnecessary	More than 1 hr.	More than 1 hr.	More than 1 hr.
Sensitivity	Up to tens of ng proteins			

● Bullet CBB Stain Lite


» In one step, complete staining in 15 minutes

No need for gel pre-staining washing

Clear staining image without the need for a destaining step

» Free from strong irritants like acetic acid and methanol

Observable staining image in the staining solution



<Conditions>


Sample : *E. coli* was prepared in Sample Buffer Solution with 2-ME(2x) for SDS-PAGE #30566-22

SDS-PAGE : Bullet PAGE Plus Precast Gel, 5-20%, 17wells (#21794-54) 400 V (constant voltage), for 10 minutes

CBB stain : Bullet CBB Stain Lite (#21964) at room temperature for 15 minutes

● CBB Stain One


- » Single bottle
- » CBB G-250 type
- » Free from acetic acid and methanol



CBB Stain One
(#04543)
Washing: 3 times x 5 mins.
Staining: 60 mins.
Destaining: Over night

● CBB Stain One Super

- » Single bottle
- » CBB R-250 type
- » Free from acetic acid and methanol



CBB Stain One Super
(#11642)
Washing: 3 times x 5 mins.
Staining: 30 mins.
Destaining: Over night

<Conditions>

Sample : Diluted Protein Markers (10x)(#29458-24) into ① 3x, ② 1x, ③ 1/3x, and ④ 1/10x, then added 2 μ L.

SDS-PAGE : Extra PAGE One Precast Gel, 5-20%, 13wells (#13063-74) 400V (constant voltage), for 30 mins.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Bullet CBB Stain Lite	R	21964-24	50 ml
		21964-95	500 ml
CBB Stain One Super (Ready-to-use)	RT	11642-31	1 L
CBB Stain One (Ready-to-use)	RT	04543-51	1 L
		04543-64	5 L
Rapid Stain CBB Kit (Coomassie R-250)	RT	30035-14	1 set (for 2 L)

Related Products

Product Name	Storage	Product No.	PKG Size
Coomassie Brilliant Blue G-250	RT	09409-42	25 g
Coomassie Brilliant Blue R-250	RT	09408-52	25 g
Ponceau S	RT	28322-72	25 g

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer


Silver Staining Kit

Silver staining method is a high sensitive method for detecting nucleic acids and proteins in polyacrylamide gel. We offer two types of silver staining kits, each having unique features for your experimental needs.


Selection of Silver Staining Kit

	Sil-Best Stain One	Sil-Best Stain-Neo
2-Dimensions	Excellent	Poor
SDS-PAGE	Good	Good
Nucleic Acid	Poor	Good
Step	12	6
Staining time	80 min.	60 min.

Comparison of Each Procedure



Comparison of Each Staining Image



● Sil-Best Stain One

Sil-Best Stain One is based on the silver staining method for protein detection. Its composition does not contain glutaraldehyde affects a result of mass spectrography.

- » **High sensitivity and low background**
- » **No glutaraldehyde**

More visible protein spot numbers than Competitors' silver staining kit




● Sil-Best Stain-Neo


Sil-Best Stain-Neo is a highly sensitive method for detecting nucleic acids and proteins in polyacrylamide gel. It is 50-100 fold more sensitive than coomassie brilliant blue and ethidium bromide.

» **Only 6 steps within up to 1 hour**

Staining Image of Protein



Staining Image of Nucleic Acid



Ordering Information

Product Name	Storage	Product No.	PKG Size
Sil-Best Stain One	R	06865-81	1 set
Sil-Best Stain-Neo for Protein and Nucleic Acid/PAGE	R	05773-11	1 set

Related Products

Product Name	Storage	Product No.	PKG Size
Dispotray-SS	RT	19386-74	40 pieces
Dispotray-S (for minigel staining)	RT	16526-82	25 pieces
Dispotray-M	RT	16551-84	20 pieces

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer













[Ultra-fast Blotting Buffer]

Bullet Semi-dry Transfer One

- » **Transfer can be completed at high speed**
- » **Efficient transfer a wide range of molecular weights**
- » **No need for gel equilibration process**
- » **Ready-to-use**

Application 1

After electrophoresis on various gels, high speed transfer was performed by using this product. EGFR, β -Actin, and Histone H3 were detected through western blotting. Proteins ranging from low molecular weight (Histone H3) to high molecular weight (EGFR) have been successfully transferred by using this product and detected quantitatively.




Gel		Bullet PAGE One Precast Gel	Extra PAGE One Precast Gel	Laemmli method gel (Hand made gel)	WIDE RANGE Gel (Hand made gel)
Blotting images	EGFR (175 kDa)				
	β -Actin (42 kDa)				
	Histone H3 (17 kDa)				

(Conditions)

- Samples : HeLa cell suspension (2 times dilution starting from 10 ug)
- SDS-PAGE : Bullet PAGE One Precast Gel, 5-15%, 17wells (#13080-44) 400 V
 Extra PAGE One Precast Gel, 5-15%, 17wells (#13062-84) 300 V
 Laemmli method gel (EGFR; 6%, β -Actin; 12%, Histone H3; 15%) 200 V
 WIDE RANGE Gel*(EGFR; 6%, β -Actin; 10%, Histone H3; 12%) 200 V
 *Casted with WIDE RANGE Gel Preparation Buffer(4x) for PAGE (#07831-94)
- Transfer : Bullet Semi-dry Transfer One (#15353-01) 25 V 10 min PVDF (poresize; 0.45 μ m)
- Blocking : Bullet Blocking One for Western Blotting (#13779) RT 5 min
- Primary antibody (EGFR) : EGF Receptor (D38B1) XP Rabbit mAb (CST #4267) 2,000x diluted
 by Signal Enhancer HIKARI for Western Blotting and ELISA (#02270-81) Solution ART 60 min
- (β -Actin) : β -Actin (C4) (Santa Cruz #sc-47778) 5,000x diluted
 by Bullet ImmunoReaction Buffer (#18439) RT 30min
- (Histone H3) : Histone H3 (D1H2) XP Rabbit mAb (CST #4499) 2,000x diluted
 by Bullet Blocking One for Western Blotting (#13779) RT 60 min
- Secondary antibody (EGFR) : Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP](Pre-adsorbed) (Novus #NB7187) 50,000x diluted
 by Signal Enhancer HIKARI for Western Blotting and ELISA (#02270-81) Solution B RT 60 min
- (β -Actin) : Goat anti-Rat, Mouse IgG (H+L) Secondary Antibody [HRP](Pre-adsorbed) (Novus #NB7574) 80,000x diluted
 by Bullet ImmunoReaction Buffer (#18439-85) RT 30min
- (Histone H3) : Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP](Pre-adsorbed) (Novus #NB7187) 30,000x diluted
 by Bullet Blocking One for Western Blotting (#13779) RT 60 min
- Detection : ChemiDoc Touch MP 3x3 binning (Bio-rad) (AUTO mode)
- (EGFR) : Chemi-Lumi One Super (#02230)
- (β -Actin) : Chemi-Lumi One L (#07880)
- (Histone H3) : Chemi-Lumi One Super(#02230)

Comparison with conventional transfer method

After electrophoresis on the Bullet PAGE Plus Precast Gel, rapid transfer (semi-dry) and conventional transfer (semi-dry and tank) were processed by this product, then detected GAPDH through western blotting. This product enables rapid transfer in just 10 minutes, achieving detection results comparable to the traditional method.

	Conditions	Blotting images
High-speed transfer	Bullet Semi-dry Transfer One(#15353-01) 25 V 10 mins Semi-dry transfer (no need for equilibration)	
Conventional transfer	Semi-dry Blotting Buffer Solution for Western Blotting 10 V for 45 min Semi-dry transfer (15 mins. equilibration performed)	
	Towbin method composition transfer buffer* 100 V tank transfer for 60 mins (15 mins equilibration performed) *25 mM Tris, 192 mM Glycine, 20%MeOH	

(Conditions)

Samples : HeLa cell suspension (2 times dilution started from 4 ug)

Transfer : Conditions listed above


Primary antibody : GAPDH Antibody (Novus #NB300-322) 10,000x diluted
by Bullet ImmunoReaction Buffer (#18439) RT 30 min

Secondary antibody : Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP](Pre-adsorbed) (Novus #NB7187) 200,000x diluted

Detection : Exposure time 3 mins

Application 2

After running on Mini-PROTEAN TGX Gels, high-speed transfer was performed by this product and transfer was confirmed by CBB staining.



[Conditions]

Sample:

① Precision Plus Protein™ All Blue Prestained Protein Standards (Bio-rad #1610373)

② 293T Cell Suspension

③ 293T Cell Suspension (2x diluted by ②)

SDS-PAGE : 4-20% Mini-PROTEAN TGX Gels (Bio-rad #4561095)

Transfer : Bullet Semi-dry Transfer One (#15353-01) 25 V for 10 min

CBB Staining : Bullet CBB Stain One(Ready To Use) (#13542)

Data provided by a University researcher.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Bullet Semi-dry Transfer One	RT	15353-01	1 L

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

[Blocking Buffer]

Blocking One and Blocking One P

Blocking is indispensable in immunoassays to block non-specific binding reactions. As Blocking One contains high molecular weight compounds, casein and bovine serum protein, it is superior to conventional blocking solutions.


Blocking One-P is an exclusive blocking solution, free of phosphate group and endogenous phosphatase for phosphorylated protein detection. The performance is superior compared with conventional blocking solutions such as 1% BSA.

The preservative in our blocking solutions do not affect the enzyme activity of peroxidase (POD) or alkaline phosphatase (ALP). Only simple refrigerator storage is necessary, even after opening the bottle.

» In many assays, a reduction of incubation time for blocking can be achieved

» Simple storage in a refrigerator even after opening the bottle

Comparison of Blocking Efficiency




The figure left shows the relationship between the reaction time and the effect of blocking in microplate assay.


In comparison with other products, Blocking One offers the strongest blocking efficiency along with faster blocking treatment time.

Western Blotting

Blocking One



Blocking One-P



Comparison with Conventional Blocking Agents

	Composition	Treatment Time	Blocking Efficacy	Phospho-specific antibody applications
Bullet Blocking One	- Amphiphilic compound - High molecular weight compounds - BSA - Casein - refer to p75	5 min.	+++	+
Blocking One	- High molecular weight compounds - BSA - Casein	20-30 min.	+++	+
Blocking One-P	- High molecular weight compounds - BSA	20-30 min.	+++	+++
Skim milk	- Casein	1 hour	+	-
1% BSA	- BSA	1 hour	+	++

Ordering Information

Product Name	Storage	Product No.	PKG Size
Blocking One	R	03953-95	500 ml
Blocking One-P	R	05999-84	200 ml




[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

[Ultra-fast Blocking Buffer]

Bullet Blocking One for Western Blotting

- » **Fast blocking in 5 mins**
- » **Ready-to-use**
- » **Simple storage in a refrigerator even after opening the bottle**

Comparison Data with Blocking One (our standard blocking reagent) and 5% skim milk














Blocking Reagents	Bullet Blocking One	Blocking One	5% Skim Milk
Blocking Time	5 min	30 min	60 min
Blotting Image			

(Conditions)

Sample : 20 µg of HeLa cell extract, 5 serial two-fold dilution series
 SDS-PAGE : Bullet PAGE One 5-15% (Product No. 13080) with SDS Running Buffer (Product No. 30329) at 400 V for 10 min.
 Blotting : Semi-dry Blotting Buffer Solution (Product No. 30650) at 10 V for 30 min
 Washing : 0.1% t-TBS (Product No. 12750)
 Blocking : Refer to the figure above
 1st antibody : Anti-Vimentin (C-20) (Rabbit) (Product No. SC-7557-R) diluted 1:2,000, 1 hr. at RT
 2nd antibody : Anti-Rabbit IgG-HRP (Product No. SC-2004) diluted 1:100,000, 1 hr. at RT
 Detection : Chemi-Lumi One (Product No. 11644)
 Detector : LAS-3000 (High mode), 15 min. exposure time

Comparison Data: Blocking efficiency in 5 minutes

The original Blocking One, the competitors' ready-to-use blocking reagents and the conventional blocking reagents did not show enough blocking efficiency in 5 mins, while Bullet Blocking One performed well.

5 min. Blocking Time	Blocking Reagents	Manufacturers' Recommended Blocking Time
	Bullet Blocking One	
	Blocking One	 30 min.
	Company A	 60 min.
	Company B	 30 min.
	Company C	 60 min.
	5% Skim Milk in 0.05 Tween® 20-TBS	 60 min.
	3% BSA in 0.05% Tween® 20-TBS	 60 min.


(Conditions)

PVDF membranes dot-blotted with mouse serum were washed with TBS. Blocking was performed using each reagent above. Anti-mouse IgG (Product No. SC-2005) (1:5,000 in 0.01% t-TBS) was applied, and the membranes were washed with 0.05% t-TBS. After reaction with Chemi-Lumi One Super (Product No. 02230), detection was performed using LAS-3000 (High mode) with 90 sec. exposure time.


Comparison of Blocking Time and Efficiency with 5% Skim Milk

Bullet Blocking One performed better than 5% skim milk in blocking time and efficiency.

(Analysis of Background Noise on Dot Blotting Data)



(Dot Blotting Data)



(Conditions)

PVDF membranes dot-blotted with mouse serum were washed with TBS. Blocking was performed using each reagent above. Anti-mouse IgG (Product No. SC-2005) (1:5,000 in 0.01% t-TBS) was applied, and the membranes were washed with 0.05% t-TBS. After reaction with Chemi-Lumi One Super (Product No. 02230), detection was performed using LAS-3000 (High mode) with 90 sec. exposure time. Data analysis was done with Multi Gauge.

Use as antibody diluent

Diluting antibody with Bullet Blocking One resulted in less non-specific binding compared to using t-TBS as diluent.

Antibody diluent	Bullet Blocking One (Undiluted)	0.1% t-TBS
Blotting Image		

(Conditions)

Sample : 10 µg of HeLa cell extraction, 3 serial two-fold dilution series
 SDS-PAGE : Bullet PAGE One 5-15% (Product No. 13080) with SDS Running Buffer (Product No. 30329) at 400 V for 12 min.
 Blotting : Semi-dry Blotting Buffer Solution (Product No. 30650) at 10 V for 30 min.
 Washing : 0.1% t-TBS (Product No. 12750)
 Blocking : Bullet Blocking One, 5 min.
 1st antibody : Anti-Cox4 (D-20) (Goat) (Product No. SC-69359) diluted 1:500, 1 hr. at RT
 2nd antibody : Anti-Goat IgG-HRP (Product No. SC-2350) diluted 1:5,000, 1 hr. at RT
 Detection : Chemi-Lumi One Super (Product No. 02230), 1 min. reaction time
 Detector : LAS-3000 (High mode), 10 min. exposure time

(Please note)

For antibody dilution, use Bullet Blocking One undiluted, or dilute up to 20x with TBS or PBS containing 0.05-0.1% detergent, such as Tween 20. Since the appropriate dilution ratio depends on antibody conditions, such as type and concentration, pretest is required.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Bullet Blocking One for Western Blotting	R	13779-56	50 ml
		13779-14	200 ml
		13779-01	1 L











[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

[Ultra-fast Antibody Dilution]

Bullet ImmunoReaction Buffer

- » **Saving-time** : Cut in half antigen-antibody reaction time
- » **Simple** : Replace tTBS, tPBS, skim milk or BSA with Bullet ImmunoReaction Buffer
- » **Ready-to-use** : Do not need to dilute

Reference data: Comparison of sensitivity related to Ab-Ag reaction time with 0.1% tTBS







Primary Ab incubation time	15 minutes	30 minutes	60 minutes	120 minutes	Overnight at 4°C
Secondary Ab incubation time	15 minutes	30 minutes	60 minutes	120 minutes	60 minutes
Bullet ImmunoReaction Buffer					
0.1%-tTBS					

(Conditions)

Sample : Hela cell extract; (left) 2 µg, (right) 0.4 µg
 Blocking : Incubated for 5 minutes with Bullet Blocking One for Western Blotting manufactured by Nacalai, #13779
 Primary antibody : β-Actin antibody manufactured by Santa Cruz, #sc-47778, x 1,000 dilution
 Secondary antibody : Goat anti-mouse IgG(H+L) secondary antibody manufactured by Novous, NB7574, x 40,000 dilution
 Detection : Chemi-Lumi One Super manufactured by Nacalai, #02230, 5-minute exposure time
 Detector : LAS-300 High mode

Transition to saving-time protocol

If a signal enhancer reagent, such as HIKARI, is use, speeding up the process with this product is not recommended, as it may decrease sensitivity and increase non-specific reactions.

Antibody diluent	ImmunoReaction Buffer	tTBS	HIKARI →refer to p79	Company A	Company B	Company C
Western blotting						
Primary antibody	30 minutes	60 minutes				
	EGF Receptor(D38B1) XP® Rabbit mAb(CST #4267), x 1,000 dilution XP is a trademark by Cell Signaling Technology, Inc.					
Secondary antibody	30 minutes	60 minutes				
	Goat anti-Rabbit IgG(H+L) Secondary Antibody[HRP](Pre-adsorbed)(Novus #NB7187), x 50,000 dilution					

(Conditions)

Sample : Hela cell extract, (left) 10 µg and (right) 2 µg
 Blocking : Incubated for 5 minutes with Bullet Blocking One
 Detection : Chemi-Lumi One Super, #02230; 3-minute exposure time
 Detector : LAS-300 High mode

Ordering Information

Product Name	Storage	Product No.	PKG Size
Bullet ImmunoReaction Buffer	R	18439-85	500 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

[High-performance Antibody Dilution]


Signal Enhancer HIKARI for Western blotting and ELISA

Dilute your antibodies with Signal Enhancer HIKARI instead of conventional diluents such as PBS-t or TBS-t before performing your next western blotting detection protocol and witness a remarkable increase in the ability to detect the protein of interest and to eliminate undesired background. Signal Enhancer HIKARI was developed to resolve the problems of low sensitivity and high background often encountered during procedures such as western blotting and ELISA.


- » Enhances antigen-antibody reactions
- » Removes background
- » Works with any substrate and membrane
- » Ready-to-use reagent



Improved Detection Sensitivity



Improved Specificity



Referenes

1. Feng-Ming Yang *et al.* *FEBS* **276**, 425-436 (2009)
2. Jian-Bin Wang *et al.* *The Journal of Cell Science* **122**(12), 2024-2033 (2009)
3. Chunwei Huang *et al.* *Reproductive Toxicology* **27**, 103-110 (2009)
4. Sawako Yamashiro *et al.* *The Journal of Cell Science* **121** (Pt 23), 3867-3877 (2008)

Ordering Information

Product Name	Storage	Product No.	PKG Size
Signal Enhancer HIKARI for Western Blotting and ELISA			
Kit contents: Solution A for Primary Antibody	R	02267-41	1 set (50 ml each)
Solution B for Secondary Antibody		02270-81	1 set (250 ml each)
Signal Enhancer HIKARI for Western Blotting and ELISA Solution A	R	02272-74	250 ml
Signal Enhancer HIKARI for Western Blotting and ELISA Solution B	R	02297-64	250 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

HRP-conjugated secondary antibody

» Affinity-purified form







» Cross-absorbed form*

*Anti-Mouse IgG: Cross-absorbed with Human IgG, Rat IgG, and Rabbit IgG

Anti-Rabbit IgG: Cross-absorbed with Human IgG, Rat IgG, and Mouse IgG

Application data

We detected target proteins in liver cells derived from induced pluripotent stem (iPS) cells, as well as in cells treated for Protein Kinase C activation. These proteins are related to cellular differentiation and phosphorylation processes. We employed three different chemiluminescent reagents with varying sensitivities to carry out the detection.

Chemiluminescent reagent (Recommended dilution rate for secondary antibodies)	Actual dilution rate	iPS cell-derived liver cells	PKC-activated HEK293 cells
Chemi Lumi One Ultra (1:50,000-1:500,000)	100,000x		
Chemi Lumi One Super (1:10,000-1:100,000)	20,000x		
Chemi Lumi One L (1:2,500-1:25,000)	5,000x		

Blue: Anti-Mouse IgG(#21860) Pink: Anti-Rabbit IgG(#21858)

(Conditions)

Sample: ① Control (l iPS cells 10 µg) ② iPS cell-derived liver cells 10 µg ③ Control (l HEK293 cells 10 µg) ④ PKC-activated HEK293 cells 10 µg

Primary antibody: Diluted with Bullet ImmunoReaction Buffer (#18439-85) and incubated for 30 mins at room temperature.

[αAT] (Proteintech #66135-1-Ig), [Oct4] (Abcam #ab19857), [β-Actin] (MBL #M177-3), [pERK] CST #4370S

[ERK] (Santa Cruz #sc-514302), [GAPDH] (Novus Biologicals #NB300-322)

Secondary antibody: Diluted with Bullet ImmunoReaction Buffer (#18439-85) and incubated for 30 mins at room temperature.

Comparison of detection sensitivity by western blotting

We detected samples that were serially diluted two-fold, from 16 µg/well using Anti-Mouse IgG and Anti-Rabbit IgG antibodies sourced from various companies at a 20,000-fold dilution. Our product demonstrated equal or higher sensitivity compared to products from other companies.

	This product	Company A	Company B
Anti-Mouse IgG (Primary antibody: Anti-β-Actin)			
Anti-Rabbit IgG (Primary antibody: Anti-PCNA)			

(Conditions)

Reference (<https://www.nacalai.co.jp/products/entry/d005021.html>)

Ordering Information

Product Name	Storage	Product No.	PKG Size
Anti-Mouse IgG(Goat), HRP-conjugated, Pre-absorbed	R	21860-74	100 µL
		21860-61	1 ml
Anti-Rabbit IgG(Goat), HRP-conjugated, Pre-absorbed	R	21858-24	100 µL
		21858-11	1 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

[Detection]

Chemiluminescent Western Blotting Substrates

Chemi-Lumi One is a series of high sensitive luminol-based chemiluminescence assay Kits for western blotting. Three types of chemiluminescent substrates are available for western blotting detection with horseradish peroxidase enzyme (HRP).

Product Name	Chemi-Lumi One L	Chemi-Lumi One Super	Chemi-Lumi One Ultra
Product No.	07880	02230	11644
Lower Detection Limit	Low-picogram	Mid-femtogram	Low-femtogram
Required Working Solution	Approx. 0.125 ml / cm ²	Approx. 0.1 ml / cm ²	Approx. 0.1 ml / cm ²
Suggested Antibody Dilution Ratio	Primary: 1:1,000-1:5,000 Secondary: 1:20,000-1:100,000	Primary: 1:1,000-1:20,000 Secondary: 1:20,000-1:200,000	Primary: 1:5,000-1:100,000 Secondary: 1:100,000-1:500,000
Reaction Period	1 min.	1 min.	5 min.
Comparable to	ECL (Cytiva) ECL Start (Cytiva) SuperSignal Pico (Thermo)	ECL Prime (Cytiva) SuperSignal Dura (Thermo) SuperSignal PLUS (Thermo) Clarity (Bio-Rad)	ECL Select (Cytiva) SuperSignal Femto (Thermo) Clarity Max (Bio-Rad)
Sensitivity	<div> <div>General Use</div> <div>High Sensitivity</div> <div>Ultra high Sensitivity</div> </div>		
<Conditions>	<div> <div>Conc. of IgG HRP-linked</div> <div> <div>Picogram</div> <div>Femtogram</div> </div> <div> <div>900</div><div>300</div><div>100</div><div>33.3</div><div>11.1</div><div>3.7</div><div>1.2</div><div>412</div><div>137</div><div>46</div><div>15</div><div>3</div> </div> </div>		
Antigen	: Anti-Mouse IgG (Goat), HRP Conjugated (Santa Cruz, sc-2005)		
Detection	: L (1 min.) Super (1 min.) Ultra (5 min.)		
Detector	: LAS-3000 Super mode (Analyze 3 min. later after reaction with each substrate)		
Ex. time	: 30 min.		

● Chemi-Lumi One L

- » Suitable for optimization of target proteins
- » Reasonable price
- » Detect wide range of protein concentration



Comparison of sensitivity with competitors

Chemi-Lumi One L offers similar sensitivity to competitors' and higher sensitivity than Company A's products.

β-Actin(ng)	50	25	12.5	6.25	3.13	<Conditions>
Chemi-Lumi One L						Gel : 10% PAGE gel Wash : 0.1% t-TBS(1x), pH7.4 Blocking : Blocking One 1st Ab : Anti-β-Actin Mouse Monol Antibody, (Santa Cruze sc-47778), 1:1,000 2nd Ab : Anti-Mouse IgG(Goat)-HRP, (Santa Cruze sc-2005), 1:20,000 Detection : 1min. Detector : LAS-3000 High mode Ex. time : 5 min.
Company A						
Company B						
Company C						


● Chemi-Lumi One Super

- » Extendable of exposure time
- » Detects proteins at mid-femtogram level with low background
- » Rapid substrate processing of blot



Comparison of sensitivity with competitors

Chemi-Lumi One Super offers the highest sensitivity out of competitors' substrates even though its exposure time is 1 minute, while others require 5 minutes.




● Chemi-Lumi One Ultra

- » The most sensitive in Chemi-Lumi One series
- » Longer signal duration
- » Wider range of experimental conditions due to low background



Comparison of sensitivity with competitors

Western blot of identical transferrin samples. The membranes were incubated with substrate that was prepared according to the manufacturers' instructions. The membranes were exposed to film for 2 minutes.



<Conditions>

Gel : 10% acrylamidegel
 Wash : 0.1% t-TBS(x1), pH7.4
 Blocking : Blocking One
 1st Ab : Anti-β-Actin Mouse Monoclonal Antibody, (Santa Cruze sc-47778), 1:300
 2nd Ab : Anti-Mouse IgG (Goat), HRP Conjugated, (Santa Cruze sc-2005), 1:1,000

Detection period : 5 min.
 Detector : LAS-3000 High mode
 Expose time : 3 min.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Chemi-Lumi One L, Luminol 250 ml and Peroxide 250 ml Sufficient substrate for 4,000 cm ² of blotting membrane	R	07880-70	1 KIT
Chemi-Lumi One Super, Luminol 50 ml and Peroxide 50 ml Sufficient substrate for 1,000 cm ² of blotting membrane	R	02230-30	1 KIT
Chemi-Lumi One Ultra, Luminol 50 ml and Peroxide 50 ml Sufficient substrate for 1,000 cm ² of blotting membrane	RT	11644-40	1 KIT

[Storage] RT = Room Temperature, R = Refrigerator, F = Freeze

Colorimetric Western Blotting Substrates








Colorimetric detection enables detection of a target protein on a membrane by a simple procedure without usage of detection equipment. Depending on the enzyme type conjugated to the antibody, some detection kits are available.

● TMB Solution for western blotting

TMB (tetramethyl benzidine) solution is used with horseradish peroxidase (HRP)-based western blotting procedure, but not applicable for ELISA or immunohistochemistry.

- » **Convenience** : No CCD-camera-equipped imager or X-ray film needed
- » **Sensitivity** : Low picogram level
- » **Safety** : Does not contain DAB or o-Dianisidine
- » **Ready-to-use** : Does not need to dilute

Reference data: Comparison with other picogram-level chemiluminescence kits and colorimetric substrates

TMB Solution for western blotting; incubation time, 10 minutes	
	TMB Solution for Western Blotting
Chemiluminescence Kits / exposure time, 10 minutes	
	Chemi-Lumi One L (#07880)
	Company A
	Company B
Other colorimetric substrates / incubation time, 10 minutes	
	Peroxidase Stain Kit (#26652-70)
	Peroxidase Stain DAB Kit (#25985-50)
	Peroxidase Stain DAB Kit +Metal Enhancer for DAB Stain (#07388-24)

Sample : HeLa cell extraction, 2 times dilution started from 12 ug
 Primary antibody : Anti-β-actin mAb manufactured by MBL, #177-3, x 5,000 dilution
 Secondary antibody : Anti-mouse IgG, HRP-linked whole Ab Sheep manufactured by GE, #NA931, x 10,000 dilution
 Detector for chemiluminescence : LAS-3000 Super mode


● BCIP-NBT Solution (Ready-To-Use)

A ready-to-use solution of chromogenic substrate of alkaline phosphatase (AP). It can be used in a wide range of applications, such as detection of alkaline phosphatase (AP) bound by antigen-antibody reaction and endogenous alkaline phosphatase (AP) of stem cells and osteoblasts, in blotting and tissue or cell staining.

- » **Ready-to-use**
- » **Can be used in wide range of applications such as immunochemical staining or endogenous AP staining**
- » **Can be used in situ hybridization (ISH), as it is tested for DNase and RNase**

Comparison: Detection sensitivity by dot blotting

Compare with other brands, BCIP-NBT Solution has equal or better sensitivity.



Ordering Information

Product Name	Storage	Product No.	PKG Size
TMB Solution for Western Blotting	R	18186-24	200 ml
BCIP-NBT Solution (Ready To Use)	R	19880-84	100 ml

Related products

Product Name	Storage	Product No.	PKG Size
Peroxidase Stain DAB Kit (Brown Stain)	R	25985-50	1 KIT
Metal Enhancer for DAB Stain	RT	07388-24	100 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer


WB Stripping Solution

WB Stripping Solution removes antibodies from blotting membrane, enabling subsequent detections with different antibodies on the very same blot. After the first antigen-antibody reaction and following chemiluminescent visualization, the antibodies can be removed by the WB Stripping Solution. A second antigen-antibody reaction can be conducted on the same blot. The same blot can be probed 2-5 times if chemiluminescent detection is employed.

- » **No heating** : **Reaction at room temperature**
- » **No odor** : **Does not contain 2-mercaptoethanol**
- » **Fast** : **Stripping time 5-15 minutes**
- » **Ready-to-use** : **One solution in one bottle**



Applications



First antigen-antibody reaction

- Blocking : Blocking One (Product No.: 03953-95), 30 min
- Wash : t-Tris Buffered Saline
- Primary ab : Anti-Paxillin (mouse IgG)
- Secondary ab : Anti-mouse IgG-POD
- Detection : Chemiluminescence Detection Kit (commercially available product)


Stripping

Condition: RT, 15 min for conventional protocol

Second, different antigen-antibody reaction

- Blocking : Blocking One (Product No.: 03953-95), 30 min
- Wash : t-Tris Buffered Saline
- Primary ab : Anti-Vinculin (mouse IgG) / Anti-Actin (mouse IgG)
- Secondary ab : Anti-mouse IgG-POD
- Detection : Chemi-Lumi One

Comparison of WB Stripping Solution and WB Stripping Solution Strong



Apply HPR-labeled anti-GST antibody to 5000 ng, 500 ng, 50 ng, or 5 ng (as desired) of c-Myc-GST antigen on a PVDF membrane, then remove the antibody by agitating gently for 10 minutes using one of the following stripping solutions.

- a: 0.05%(v/v) t-TBS
- b: 2%(w/v) SDS, 100mM 2-Meraptoethanol
- c: **WB Stripping Solution**
- d: **WB Stripping Solution Strong**

After stripping the antibodies and washing the membrane with t-PBS for 2 min, use the chemiluminescence method to detect the HPR-labeled anti-GST antibody remaining on the membrane.

*Image "e" is a result that shows detection of the antigen with HPR-labeled anti-GST antibody on the "d". The similar result is marked with "a". Therefore, WB Stripping Solution Strong only stripped antibodies, not antigens.

Ordering Information

Product Name	Storage	Product No.	PKG Size
WB Stripping Solution	RT	05364-55	500 ml
WB Stripping Solution Strong	R	05677-65	500 ml
WB Stripping Solution Trial Set (WB Stripping Solution: 40 ml, WB Stripping Solution Strong: 40 ml)	R	05680-21	1 set

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer


Epitope Tag Antibody

A family of epitope tag antibodies available for the detection and purification of the recombinant proteins. Most of our tag antibodies are highly specific mouse and rat monoclonal antibodies.

● Anti-GFP (Rat IgG2a), Mono (GF090R)

Clone	: GF090R
Isotype	: IgG2a (Rat)
Product form	: Liquid
Immunogen	: His-GFP (full length) fusion protein
Application	: Immunohistochemistry 1:1000-1:2000
	: Western Blotting 1:1000-1:2000
	: ELISA 1:2000-1:20000


Immunohistochemistry



Sample	: Mouse brain (nerve cell)
Primary antibody	: Anti-GFP(Rat IgG2a), Mono (1:1000) RT, O/N
Secondary antibody	: Anti-Rat IgG-Cy3 (1:300) RT, 1 hr
Blocking	: 5% Normal goat serum/0.2% TritonX-100 in PBS
Fixing method	: 4% Paraformaldehyde

Data courtesy of Dr. Y. Yoshihara, RIKEN Brain Science Institute

Western Blotting




Reference

1. Nakamura, M. *et al. Molecular Vision* **16**, 425-437 (2010)
2. Nishide, K. *et al. PLoS ONE* **4**(8), e6869 (2009)
3. Nagao, M. *et al. The Journal of Cell Biology* **183**(7), 1243-1257 (2008)
4. Ono, K. *et al. Development Biology* **320**(2), 356-468 (2008)
5. Nakashiba T. *et al. Science* **319**(5867), 1260-1264 (2008)
6. Batista-Brito R *et al. The Journal of Neuroscience* **28**(15), 3966-3975 (2008)
7. Esumi S. *et al. Neuroscience Research* **60**(4), 439-451 (2008)
8. Fogarty M. *et al. The Journal of Neuroscience* **27**(41), 10935-10946 (2007)
9. Sasamura T. *et al. Development* **134**, 1347-1356 (2007)
10. Sato Y. *et al. The Journal of Neuroscience* **27**(7), 1606-1615 (2007)
11. Yamada M. *et al. Stem Cells* **25**(3), 562-570 (2007)
12. Togashi H. *et al. The Journal of Cell Biology* **174**, 141-151 (2006)
13. Ogata M. *et al. Molecular and Cell Biology* **26**(24), 9220-9231 (2006)
14. Sato Y. *et al. The Journal of Neuroscience* **25**(20), 4889-4897 (2005)


● Anti-GFP (Mouse IgG1-k), Mono (GF200)

Clone	: GF200
Isotype	: IgG1-k (Mouse)
Product form	: Liquid
Immunogen	: His-GFP (full length) fusion protein
Application	: Western Blotting 1:1000-1:2000
	: ELISA 1:2000-1:20000

Western Blotting



Western Blotting



● Anti-c-Myc (Mouse IgG1-k), Mono (MC045)

Clone	: MC045	
Isotype	: IgG1-k (Mouse)	
Product form	: Liquid	
Immunogen	: c-Myc synthetic peptide [EQKLISEEDL] conjugated with KLH	
Application	Western Blotting	1:1000-1:2000
	Immunoprecipitation	1:400-1:1000
	Immunocytochemistry	1:400-1:1000
	ELISA	1:2000-1:20000

Immunocytochemistry



Sample	SH-SY5Y cell in which c-Myc-tagged TCP1β is expressed	
Primary antibody	: Anti-c-Myc (Mouse IgG1-κ) (2.5 μg/ml)	
	Anti-Protein X	
Secondary antibody	: Anti-Fluor 546-conjugated antibody (1:400)	
	Anti-Fluor 488-conjugated antibody (1:400)	

Data courtesy of RIKEN Brain Science Institute

● Anti-6xHis (Mouse IgG1a-k), Mono (HI192)

Clone	: HI192	
Isotype	: IgG1-k (Mouse)	
Product form	: Liquid	
Immunogen	: 6xHis synthetic peptide [HHHHHH] conjugated with KLH	
Application	Western Blotting	1:1000-1:2000
	ELISA	1:2000-1:20000

Western Blotting

A Western blot image showing a single band at approximately 66 kDa. The molecular weight markers on the left are 199, 124, 98, 68, and 48 kDa. The band is located between the 68 and 98 kDa markers, closer to the 68 kDa marker.

Sample:
Lysate from Sf9 cells in which His-tagged
Drosophila PTEN is expressed

Filter	: FluoroTrans [PALL]
Blocking	: 5% non-fat dry milk/PBS (30 min)
Primary antibody	: Anti-6xHis (Mouse IgG1 κ , Monoclonal (HI192)
Wash	: 0.25% Tween-20/TBS (10 min x 3)
Secondary antibody	: Anti-Mouse IgG-HRP (1:1000) /2% BSA/0.25% Tween-20 /TBS (30 min)
Wash	: 0.25% Tween-20/TBS (10 min x 5)
Detection	: Luminol Reagent (Santa Cruz)

*Data courtesy of Dr. A. Maehama, The Tokyo Metropolitan
Institute of Medical Science (RINSHOKEN)*

Ordering Information

Product Name	Application	Storage	Product No.	PKG Size
Anti-c-Myc (Mouse IgG1-k), Monoclonal (MC045), AS	WB, IP, ICC, ELISA	R	04362-76	50 μg
			04362-34	200 μg
Anti-c-Myc (Mouse IgG1-κ), Monoclonal (MC045), AS, Agarose Conjugate	IP	R	04145-55	500 μg
Anti-GFP (Mouse IgG1-k), Monoclonal (GF200), AS	WB, ELISA	R	04363-66	50 μg
			04363-24	200 μg
Anti-GFP (Rat IgG2a), Monoclonal (GF090R), CC	WB, IHC ELISA	R	04404-26	50 μg
			04404-84	200 μg
Anti-GFP (Rat IgG2a), Monoclonal(GF090R), CC, Agarose Conjugate	IP	R	06083-05	500 μg
Anti-6xHis (Mouse IgG1-k), Monoclonal (HI192), AS	WB, ELISA	R	04428-26	50 μg
			04428-84	200 μg

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

4% - Paraformaldehyde Phosphate Buffer Solution

We offer a 10% formalin neutral buffer solution, which contains about 4% formaldehyde, designed for use as a general fixation buffer in histological specimen preparations. Since this product is based on commonly available formalin, methanol is used as a stabilizer.


If a methanol free formalin solution is required, paraformaldehyde is used instead of formaldehyde. However, paraformaldehyde is extremely toxic and can cause injury if scattered. To deal with this hazard, additional work, such as making the solution alkaline with heating when dissolving the paraformaldehyde, is required. Our product is available in two volume types: 500 ml and a 5 x 10 ml package set.



- » **Small unit volume**
- » **Enables to immerse histological specimens directly into the solution (5 x 10 ml)**
- » **Low cost for waste**
- » **Ready-to-use**
- » **Storable in refrigerator**


Usage examples

Hematoxylin-eosin staining



Sample : Mouse small intestine
 Fixation method : Immersion fixation with this product (over night at 4°C)
 Deparaffinization : Limonene and ethanol

Immunohistological staining



Sample : Mouse small intestine
 Fixation method : Immersion fixation with this product (overnight at 4°C)
 Deparaffinization : Limonene and ethanol
 Primary antibody : Anti-PCNA (FL-261) (rabbit)
 Staining : Peroxidase Stain DAB Kit (brown stain) (Product No. 25985-50)

Ordering Information

Product Name	Storage	Product No.	PKG Size
4% - Paraformaldehyde Phosphate Buffer Solution	R	09154-14	5 x 10 ml
		09154-85	500 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

HistoVT One (10x, pH 7.0)


HistoVT One is an antigen retrieval solution for immunohistochemistry and *in situ* hybridization. This product can unmask antigenic sites without damage to antigen from formalin-fixed, frozen or paraffin-embedded tissue sections.

- » Enhancing antigen-antibody reaction
- » Usable with frozen or paraffin-embedded tissue section
- » High reproducibility




Application 1

Formalin-fixed, paraffin-embedded tissue sections



Application 2

Frozen sections



Data courtesy of RIKEN Brain Science Institute, Brain Development Research Group

Ordering Information

	Product Name	Storage	Product No.	PKG Size
	HistoVT One (10x, pH 7.0)	RT	06380-05	500 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer


Blocking One Histo

Blocking One Histo is a blocking solution to prevent non-specific binding of antibodies in immunohistochemistry (IHC). The product is designed for immunohistochemistry application based on Blocking One (refer to Western Blotting Section).

- » Eye-drop bottle
- » Can be used for immunofluorescence staining
- » The preservative does not affect the activity of alkaline phosphatase (AP) or horseradish peroxidase (HRP)



Comparison of blocking efficiency with 10% goat serum (Immunofluorescence)




Sample : Mouse small intestine (Paraffin-embedded section)
Antigen Retrieval : Histo VT One, 90°C, 20 min.
Primary antibody : Anti-Vimentin rabbit polyclonal antibody (Santa Cruz: #sc-7557R)
Secondary antibody : CF™ 488A Goat anti-Rabbit IgG(H+L), F(ab')₂ Fragment (Biotium: #20013)

Blocking One Histo (10 min.) 10% Goat Serum (10 min.)

In both panels, mouse small intestine tissue section was stained with secondary antibody conjugated with CF™ 488A (green) and counter stained with DAPI (blue). In the panel B with 10% Goat Serum, the stained white arrow along the lines of shape of small intestine show non-specific staining. Blocking One Histo is more effective at reducing non-specific background staining than normal serum.

Applications



A: Mouse small intestine (PCNA) x5
B: Mouse epididymis (Vimentin) x25
C: Mouse brain (GluR) x100

Antigen Retrieval : Histo VT One, Room temp., 10 min.
Primary antibody : A: Anti-PCNA rabbit pAb (Santa Cruz: #sc-7907)
 B: Anti-Vimentin rabbit pAb (Santa Cruz: #sc-7557R)
 C: Anti-GluR-1 goat pAb (Santa Cruz: #sc-7608)
Secondary antibody : A: Goat anti-rabbit IgG (H+L), biotin conjugated (Vector, #BA-1000)
 B: Goat anti-rabbit IgG (H+L), biotin conjugated (Vector, #BA-1000)
 C: Bovine anti-goat IgG (H+L), biotin conjugated (Santa Cruz: #sc-2347)
Detection : Streptavidin Biotin Complex Peroxidase Kit (Product No. 30462)
 Peroxidase Stain DAB Kit (Brown Stain) (Product No. 25985)

Blocking treatment of each tissue section had been performed by Blocking One Histo. Mouse small intestine (panel A) was stained with anti-PCNA and DAB (3,3'-Diamino Benzidine) to stain nuclear (black arrow), Mouse epididymis (panel B) was stained with anti-Vimentin and DAB to stain muscle (black arrow), Mouse brain (panel C) was stained with anti-GluR and DAB to stain membrane proteins (black arrow) and counter stained with hematoxylin.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Blocking One Histo	R	06349-64	50 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer


Signal Enhancer HIKARI for Immunostain

Signal Enhancer HIKARI for Immunostain was developed to resolve the problems of low sensitivity and high background often encountered during immunostain procedures such as immunohistochemistry (IHC) and immunocytochemistry. Dilute your antibodies with Signal Enhancer HIKARI for Immunostain instead of conventional diluents such as PBS or TBS before performing your next IHC experiment and witness a remarkable increase in the ability to detect the protein of interest and to eliminate unwanted background.

- » **Enhances signals**
- » **Reduces background**
- » **Ready-to-use reagent**
- » **Works with any detection system**

* The kit can also be used in combination with sensitizing systems such as the ABC or polymer complex method.

Applications




Ordering Information

Product Name	Storage	Product No.	PKG Size
Signal Enhancer HIKARI for Immunostain Trial Set	R	02363-71	1 set (5 ml each)
Signal Enhancer HIKARI for Immunostain Solution A	R	02373-54	20 ml
Signal Enhancer HIKARI for Immunostain Solution B	R	02375-34	20 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Anti-GFP (Rat IgG2a), Monoclonal (GF090R)

- » Immunohistochemical application
- » Rat monoclonal antibody



Sample : Mouse brain (nerve cell)
 Primary antibody : Anti-GFP(Rat IgG2a), Mono (1:1000) RT, O/N
 Secondary antibody : Anti-Rat IgG-Cy3 (1:300) RT, 1 hr
 Blocking : 5% Normal goat serum/0.2% TritonX-100 in PBS
 Fixing method : 4% Paraformaldehyde

Data courtesy of Dr. Y. Yoshihara, RIKEN Brain Science Institute

References for Immunostaining

1. Takeshi Sasamura *et al. Development* **134**, 1347-1356 (2007)
2. Takashi Inoue *et al. The Journal of Neuroscience*, **27**(20), 5461-5473 (2007)
3. Anoop Kumar G. Velikkakath *et al. Mol Biol Cell*. **23**(5), 896-909 (2012)
4. Eisuke Itakura *et al. Molecular Biology of the Cell*, **19**, 5360-5372,
5. Keith N. Brown, *et al. Science*, **334**, 480 (2011)
6. Anna N. Rubin *et al. The Journal of Neuroscience*, **30**(36), 12050-12062 (2010)
7. Shinsuke Nakao *et al. J Cell Biol.*, **182**(2), 395-410 (2008)
8. Maiko Ogata *et al. Mol Cell Biol.*, **26**(24), 9220-9231 (2006)
9. Matthew Fogarty *et al. The Journal of Neuroscience*, **27**(41), 10935-10946 (2007)
10. Takuya Sato *et al. Nature Communications*, **2** (472)
11. Toshiaki Nakashiba *et al. Science*, **319**(5867), 1260-4 (2008)
12. Hiromi Takanaga *et al. Stem Cells*, **27**(1), 165-74 (2009)
13. Akinori Yamasaki *et al. Mol. Biol. Cell*, **17** (11), 4876-4887 (2006)
14. Natsumi Ageta-Ishihara *et al. The Journal of Neuroscience*, **29**(43), 13720-13729 (2009)
15. Naoyuki Asada *et al. Journal of Neuroscience*, **30**(26), 8852-8865 (2010)
16. Shizue Ohsawa *et al. Dev Cell*, **20**(3), 315-28 (2011).

Ordering Information

For more informations about Epitope Tag Antibody, please refer to p87.

Product Name	Storage	Product No.	PKG Size
Anti-GFP(Rat IgG2a), Monoclonal(GF090R), CC	R	04404-26	50 µg
Anti-GFP(Rat IgG2a), Monoclonal(GF090R), CC, Agarose Conjugate	R	06083-05	500 µg

Streptavidin Biotin Complex Peroxidase Kit

Streptavidin Biotin Complex Peroxidase Kit includes reagents for the "Avidin-Biotin Complex, ABC technique", a highly sensitive method for immunoblotting, immunohistochemistry, ELISA and *in situ* hybridization.



Ordering Information

Product Name	Storage	Product No.	PKG Size
Streptavidin Biotin Complex Peroxidase Kit	R	30462-30	1 KIT

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Nucleic Acid Isolation
/ Electrophoresis

Cell Culture

Cell Extraction
/ Protein Assay

Protein Purification

Protein Electrophoresis

Western Blotting

Immunohistochemistry

High Sensitivity Peroxidase DAB Stain

● Peroxidase DAB Stain Kit and Metal Enhancer

Peroxidase Stain DAB Kit (Brown Stain) is used to detect horseradish peroxidase (HRP) activity and stain them brown in immunoblotting, immunohistochemistry and *in situ* hybridization.

Metal Enhancer for DAB Stain (Product No. 07388-24) is used to stain peroxidase purplish gray with Peroxidase Stain DAB Kit (Brown Stain)(Product No. 25985-50) in immunoblotting, immunohistochemistry and *in situ* hybridization. The sensitivity of Metal Enhancer for DAB Stain used with Peroxidase Stain DAB Kit (Brown Stain) is about two times higher than the product with Peroxidase Stain DAB Kit (Brown Stain) alone.

- » **High sensitivity**
- » **Metal Enhancer for DAB Stain stains peroxidase purplish gray**
(Just change the solution mix from water to Metal Enhancer for DAB)
- » **RNase, DNase free, applicable to *in situ* hybridization**
- » **Eye drop bottle**



Procedure of combination Peroxidase Stain DAB Kit and Metal Enhancer Solution




Change the adjusting solution of Peroxidase Stain DAB Kit (Brown Stain) from water to Metal Enhancer for DAB.

Application

Immunohistostaining of mouse small intestines with anti-PCNA antibody (Serial membranes)

The sensitivity achieved when the Peroxidase Stain DAB Kit (Brown Stain) is used alone is higher than the competitors' products. However, when used in conjunction with Metal Enhancer for DAB stain, the sensitivity of Peroxidase Stain DAB Kit (Brown Stain) is about two times higher than what can be achieved by the Peroxidase Stain DAB Kit (Brown Stain) alone.



- NACALAI TESQUE : (Gray) Peroxidase Stain DAB Kit (Brown Stain) + Metal Enhancer for DAB
(Brown) Peroxidase Stain DAB Kit (Brown Stain)
- Vector : (Gray) Kit (with attached nickel solution)
(Brown) Kit (without attached nickel solution)
- Basic method : (Gray) 0.6mg/ml DAB, 0.03% H_2O_2 , 50mM Tris-HCl Buffer pH7.6, 0.03% NiCl_2
(Brown) 0.6mg/ml DAB, 0.03% H_2O_2 , 50mM Tris-HCl Buffer pH7.6

Ordering Information

Product Name	Storage	Product No.	PKG Size
Peroxidase Stain DAB Kit (Brown Stain)	R	25985-50	1 KIT
Metal Enhancer for DAB Stain	RT	07388-24	100 ml

[Storage] RT = Room Temperature, R = Refrigerator, F = Freezer

Mounting Medium for Fluorescent Staining

Fluoro-KEEPER Antifade Reagent is a non-hardening mounting medium with a unique antifade reagent. It suppresses rapid photobleaching during fluorescence microscopy observation. The coverslipped slide with nail polish or other sealants can be stable for several weeks. There are two types of products available, with DAPI [4',6-Diamidino-2-phenylindole] and without DAPI, which counterstains nucleus blue.

- » Inhibits photobleaching of various fluorescent dyes
- » Easy to use with eye-drop bottle


Application 1 (Fluoro-KEEPER with DAPI)

Fluorescent microscopy experiments

Fluoro-KEEPER with DAPI offers nuclear staining along with mounting.


Mounting Medium : Fluoro-KEEPER with DAPI for 30 min. at room temperature protecting from light.
Microscopy : Olympus BX-50-34-FLA1

Mouse Small Intestine




Antigen retrieval : HistoVT One (#06380)
Blocking : Blocking One Histo (#06349)
1st Ab : Anti-Vimentin Rabbit Polyclonal Antibody (Santa Cruz #sc-7557R)
2nd Ab : CFTM 488A Goat Anti-Rabbit IgG (H+L), Fragment (Biotium #20013)

MC3T3-E1 Cell



Blocking : Blocking One Histo (#06349)
Rhodamine-conjugated phalloidin (Cytoskeleton #PHDR1)

MC3T3-E1 Cell




Blocking : Blocking One Histo (#06349)
1st Ab : Anti-Vimentin Rabbit Polyclonal Antibod (Santa Cruz #sc-7557R)
2nd Ab : Cy² Goat Anti-Rabbit IgG (H+L)(GENETEX #GTX26940)

Application 2 (Fluoro-KEEPER without DAPI)


Fluorescent microscopy experiments: Planarian

Fluorescence intensities are shown as percentages of initial intensities remaining during repeated frame capture up to 60 times. The images were acquired by Olympus FV10. The samples mounted in the Fluoro-KEEPER Antifade Reagent were clearly detected after 60 times of frame capture.

■ Stain Image



■ Fluorescence intensity



Nuclear

Hoechst 33342

Arrestin and G Protein β Subunit(Gβ)

1st Ab: Mouse Anti-planarian Arrestin

Mouse Anti-planarian Gβ

2nd Ab: Alexa Fluor® 488 Goat Anti-mouse IgG

Tryptophan β Hydroxylase(TPβH)

1st Ab: Rabbit Anti-planarian TPβH

2nd Ab: Alexa Fluor® 594 Goat Anti-rabbit IgG

Samples were mounted in the Fluoro-KEEPER Antifade Reagent

Data courtesy of Agata Lab, Department of Biophysics, Kyoto University

Comparison of antifade effectiveness with different fluorescent dyes

Fluoro-KEEPER Antifade Reagent offers increased resistance to photobleaching of various fluorescent dyes.

Fluorescence Dye	without DAPI		with DAPI	
	Fluoro-KEEPER	Control	Fluoro-KEEPER	Control
Hoechst 33258	98	96	—	—
Hoechst 33342	100	90	—	—
DAPI	99	93	—	—
Propidium Iodide	95	67	—	—
Fluorescein	97	49	96	49
Alexa Fluor® 488	93	86	96	91
CF™ 488	93	82	91	82
Cy® 2	99	83	98	81
Rhodamine	72	51	78	41
Alexa Fluor® 555	98	81	97	87
CF™ 555	98	85	97	85
Cy® 3	89	71	86	66

Cells stained by each fluorescent dye were mounted in Fluoro-KEEPER Antifade Reagent, 85% Glycerol containing PBS as a control. Samples were illuminated for 60 seconds. Each number indicates fluorescence intensity as percentage of initial intensity after 60 seconds exposure photobleaching.

Control Condition: 85% Glycerol-PBS w/o DAPI
85% Glycerol-PBS w/ DAPI

Fluorescent Microscopy: Olympus BX-50-34-FLA1

Exposure time: 60 seconds.

Ordering Information

Product Name	Storage	Product No.	PKG Size
Fluoro-KEEPER Antifade Reagent, Non-Hardening Type	R	12593-64	2 x 5 ml
Fluoro-KEEPER Antifade Reagent, Non-Hardening Type with DAPI	R	12745-74	2 x 5 ml

[Storage] RT = Room temperature, R = Refrigerator, F = Freezer

www.nacalai.com

NACALAI TESQUE, INC.

Nijo Karasuma, Nakagyo-ku, Kyoto 604-0855 JAPAN

TEL : +81 (0)75 251 1730

E-mail : info.intl@nacalai.com

www.nacalai.com

ATTENTION:

Nacalai Tesque makes no representation or warranty as to whether the products and/or their uses infringe any patent of any third party, nor shall Nacalai Tesque be liable for infringement of any such patent.

Warranties and Disclaimers:

Nacalai Tesque warrants that its products shall conform to the description of such products as provided by Nacalai Tesque through its catalog, analytical data or other literature. Nacalai Tesque makes no other warranty, express or implied, as to the fitness of these products for any particular purpose. Nacalai Tesque shall not in any event be liable for incidental or consequential damages that may result from any use or failure of the products.

For more information on products and pricing, please contact your local distributor.